

Guidelines for Prehospital Fluid Resuscitation in the Injured Patient

Bryan A. Cotton, MD, MPH, Rebecca Jerome, MLIS, MPH, Bryan R. Collier, DO, Suneel Khetarpal, MD, Michelle Holevar, MD, Brian Tucker, DO, Stan Kurek, DO, Nathan T. Mowery, MD, Kamal Shah, MD, William Bromberg, MD, Oliver L. Gunter, MD, and William P. Riordan, Jr., MD; EAST Practice Parameter Workgroup for Prehospital Fluid Resuscitation

Abstract: Although the need and benefit of prehospital interventions has been controversial for quite some time, an increasing amount of evidence has stirred both sides into more frequent debate. Proponents of the traditional “scoop-and-run” technique argue that this approach allows a more timely transfer to definitive care facilities and limits unnecessary (and potentially harmful) procedures. However, advocates of the “stay-and-play” method point to improvement in survival to reach the hospital and better neurologic outcomes after brain injury. Given the lack of consensus, the Eastern Association for the Surgery of Trauma convened a Practice Management Guideline committee to answer the following questions regarding prehospital resuscitation: (1) should injured patients have vascular access attempted in the prehospital setting? (2) if so, what location is preferred for access? (3) if access is achieved, should intravenous fluids be administered? (4) if fluids are to be administered, which solution is preferred? and (5) if fluids are to be administered, what volume and rate should be infused?

Key Words: Resuscitation, Intravenous fluid, Venous access, Intraosseous, Prehospital, Field.

(*J Trauma*. 2009;67: 389–402)

STATEMENT OF THE PROBLEM

Over the past several decades, the scope of practice for emergency medical personnel has rapidly expanded.¹ Along with this, a dramatic increase in the number of prehospital procedures (especially intubation and central venous access) has been noted.^{2,3} However, this dramatic change in the prehospital approach to the injured patient

has occurred in the absence of data to support its adoption. Investigators from Los Angeles have noted no difference in survival when injured patients are transported by a private vehicle or emergency medical services (EMS) transport.^{4,5} Liberman et al.⁶ recently evaluated Canadian cities with Level I trauma centers and demonstrated no difference in outcomes between the basic life support delivered by EMS providers and the prehospital advanced life support delivered by either paramedics or physicians.

Despite a lack of evidence demonstrating a benefit to prehospital fluid resuscitation, this practice is considered to be standard of care. As well, the type of fluid, the appropriate rate of administration, and the resuscitations themselves remain unguided and unsupervised. Recently, the largest prehospital organization in the United Kingdom issued a consensus statement calling for more “restrained” and “cautious” use of crystalloids in prehospital settings.⁷ The authors argue that several prehospital hemorrhage models have demonstrated that limiting the initial resuscitation volume, before definitive care, leads to a reduction in hemorrhage.^{8,9} Unfortunately, clinical extrapolation of deliberate hypotension has, for the most part, been confined to a single prospective trial by Bickell et al.¹⁰ The investigators noted a decreased length of stay and lower mortality in patients with delayed resuscitation.

Although the need and benefit of prehospital interventions has been controversial for quite some time, an increasing amount of evidence has stirred both sides into more frequent debate. Proponents of the traditional “scoop-and-run” technique argue that this approach allows a more timely transfer to definitive care facilities and limits unnecessary (and potentially harmful) procedures. However, advocates of the “stay-and-play” method point to improvement in survival to reach the hospital and better neurologic outcomes after brain injury. Given the lack of consensus, the Eastern Association for the Surgery of Trauma convened a Practice Management Guideline (PMG) committee to answer the following questions regarding prehospital resuscitation (1): should injured patients have vascular access attempted in the prehospital setting? (2) if so, what location is preferred for access? (3) if access is achieved, should intravenous fluids be administered? (4) if fluids are to be administered, which solution is preferred? and (5) if fluids are to be administered, what volume and rate should be infused?

Submitted for publication October 24, 2008.

Accepted for publication April 5, 2009.

Copyright © 2009 by Lippincott Williams & Wilkins

From the Department of Surgery (B.A.C.), University of Texas Health Science Center, Houston, Texas; Department of Surgery (R.J., B.R.C., W.P.R.), Vanderbilt University Medical Center, Nashville, Tennessee; Department of Surgery (S.K.), Tampa General Hospital, Tampa, Florida; Department of Surgery (M.H.), Mount Sinai Hospital, Chicago, Illinois; Department of Surgery, University of Tennessee (B.T., S.K.), Knoxville Medical Center, Knoxville, Tennessee; Department of Surgery (N.T.M.), Wake Forest/Baptist Medical Center, Winston-Salem, North Carolina; Department of Surgery (K.S.), Lehigh Valley Hospital and Health Network, Allentown, Pennsylvania; Department of Surgery (W.B.), Memorial Health University Medical Center, Savannah, Georgia; Department of Surgery (O.L.G.), Washington University/Barnes Jewish Medical Center, St. Louis, Missouri.

Presented at the 21st Annual Meeting of the Eastern Association for the Surgery of Trauma, January 15–19, 2008, Jacksonville, Florida.

Address for Reprints: Bryan A. Cotton, MD, MPH, The University of Texas Health Science Center at Houston, The Center for Translational Injury Research, 6410 Fannin St., 1100.20 UPB, Houston, TX 77030; email: bryan.a.cotton@uth.tmc.edu.

DOI: 10.1097/TA.0b013e3181a8b26f

PROCESS

Identification of References

A primary computerized search of the National Library of Medicine and the National Institute of Health MEDLINE database was undertaken using the PubMed Entrez interface. This search was undertaken by a Masters of Library Sciences Faculty member at Vanderbilt University School of Medicine. All English language citations between 1982 and 2007 focusing on prehospital vascular access and fluid resuscitation of trauma patients were reviewed. This primary search query retrieved 3,300 citations:

1. fluid AND prehospital, OR out of hospital, OR field, AND resuscitation, AND injury
2. access OR vascular OR intravenous OR cannulation AND prehospital OR field OR out of hospital AND injury
3. prehospital AND injury AND dextrans/ or Sodium Chloride/ or Saline Solution, Hypertonic/ or saline OR blood substitute OR fluids
4. intraosseous AND access

Review articles, letters to the editor, case reports, and editorials were excluded. Basic science and animal studies, as well as items limited to pediatric patients and noninjured patients were also excluded. Excluding studies with the above-stated populations limited this literature query to 79 articles.

A secondary query was carried out by PMG study group members (BAC, BRC) using the OVID database version of MEDLINE with the same search terms, exclusion, and date range. The secondary search criteria yielded 92 citations after above-stated exclusions.

There were 12 citations that were missed on the primary that were detected on the secondary. Searches merged and the removal of articles that did not strictly apply to the study questions and population of interest yielded 58 articles. A total of 16 articles were reviewed by the committee and determined by the members to be technical studies (flow rates for catheters and/or delivery systems) or mixed populations of patients (medical and surgical not purely trauma). Exclusion of these left 42 total articles that formed our evidence-based review and populated Tables 1–5.

Quality of the References

The references were classified using methodology established by the Agency for Health Care Policy and Research of the U.S. Department of Health and Human Services (add reference).¹¹ Additional criteria and specifications were used for classification of articles from a tool described by Oxman et al.¹² Thus, the classifications were as follows:

- Class I: Prospective, randomized controlled trials. A total of 10 Class I articles were reviewed.
- Class II: Clinical studies prospectively collected data and retrospective analyses, which were based on clearly reliable data. Fourteen articles met criteria for Class II articles and were reviewed.
- Class III: Studies based on retrospectively collected data; includes clinical series, database or registry reviews, large series of case reviews, and expert opinion. Eighteen arti-

cles were identified as class III and underwent review. Each of the above articles was reviewed and scored by a minimum of two PMG committee members. After collection of all reviews, the prehospital Fluid Resuscitation PMG Committee convened and developed recommendations based on the following definitions:

- Level I: The recommendation is convincingly justifiable based on the available scientific information alone. One Level I guideline was supported by the literature.
- Level II: The recommendation is reasonably justifiable by available scientific evidence and strongly supported by expert opinion. A total of eight Level II guidelines were established by the literature.
- Level III: The recommendation is supported by available data but adequate scientific evidence is lacking. Seven Level III guidelines were developed.

RECOMMENDATIONS

Should vascular access be obtained in the prehospital setting? (Table 1).^{7,13–28}

- A. Level I: No level one recommendation can be made. There is insufficient data to support a Level I recommendation for placing vascular access in the prehospital setting.
- B. Level II: Placement of vascular access at the scene of injury should not be performed as it delays patient transport to definitive care, and there is no evidence to demonstrate any benefit to their placement.
- C. Level III: Placement of vascular access during transport is feasible and does not delay transport to definitive care.

If vascular access is obtained, where and how should it be placed? (Table 2).^{7,10,27,29–43}

- A. Level I: No Level I recommendation can be made. There is insufficient data to support specifically where and through which approach vascular access should be obtained in the prehospital setting of trauma.
- B. Level II: (a) If central access is necessary, the percutaneous “Seldinger” technique is recommended over traditional “cut-down” procedures as there is evidence that percutaneous techniques are quicker and have equivalent success rates. (b) The use of intraosseous access in trauma patients requiring vascular access in which intravenous access is unobtainable or has failed two attempts is recommended.
- C. Level III: Attempts at peripheral intravenous access should be limited to two attempts during prehospital transport after which, alternative methods (intraosseous, central access) should be attempted if equipment and trained personnel are available.

If vascular access is obtained, should intravenous fluids be given? (Table 3).^{7,10,27,30–34,38–42}

- A. Level I: No level one recommendation can be made. There is insufficient data to show that trauma patients benefit from prehospital fluid resuscitation.
- B. Level II: (a) Intravenous fluids should be withheld in the prehospital setting in patients with penetrating torso injuries.

TABLE 1. Should Vascular Access Be Obtained?

Author(s)	Year	Title	Class	Comments and Consensus
O'Gorman et al. ²¹	1989	Zero-time pre-hospital I.V. <i>J Trauma.</i> 1989;29:84–86.	II	Prospective observational-evaluated whether pretransport venous access was more successful than in-transport attempts. 350 patients. No statistical difference in ability to place IV prior to or during transport in nonhypotensive patient. In hypotensive patients placement in ambulance is more successful.
Jones et al. ¹⁹	1989	Prehospital intravenous line placement: a prospective study. <i>Ann Emerg Med.</i> 1989;18:244–246.	II	Prospective observational study. 97 patients; 12 patients were hypotensive or in cardiac arrest. Success rate for trauma patients was 96%. On scene success rate 91%. En route success 94%. Hypotensive 86% success rate. Average time to placement (single attempt) 2.8 min; with failed attempts, 6.3 min. En route placement recommended.
Minville et al. ²⁰	2006	Prehospital intravenous line placement assessment in the French emergency system: a prospective study. <i>Eur J Anaesthesiol.</i> 2006;23:594–597.	II	Prospective observational study. 388 patients (83% medical). Results: 76% success for first attempt. Average time to IV was 4.4 min with a 99.7% success rate. 71% of the patients received therapy through the line prehospital.
Honigman et al. ¹⁸	1990	Prehospital advanced trauma life support for penetrating cardiac wounds. <i>Ann Emerg Med.</i> 1990;19:145–150.	II	Retrospective. 70 consecutive patients with penetrating cardiac injuries (31 GSW, 39 stab wounds). Results: 10.7 min on scene, 93% had IV access gained. 30% survival. No correlation between on scene time and IV access gained. No statistical difference noted. Recommendations: Well-trained paramedics can perform procedures with short scene times and high rate of survival.
Pace et al. ²²	1999	Pace SA, Fuller FP, Dahlgren TJ. Paramedic decisions with placement of out-of-hospital intravenous lines. <i>Am J Emerg Med.</i> 1999;17:544–547.	II	Prospective observational cohort. 290 patients. Results: 57% of patients received out-of-hospital intravenous catheters with an over-treatment rate of 29% ± 5%, and an under-treatment rate of 2.4% ± 1.8%. 89% overall successful placement rate Conclusions: Lines are frequently started and not used. Paramedics exercise reasonable judgment and appropriate decision making when deciding to start an IV. 29% nonuse rate may represent acceptable trade off to achieve under-treatment rate of 2.4%.
Barrett and Guly. ¹³	2000	How long does it take to perform procedures on scene? <i>Prehospital Immed Care.</i> 2000;4:25–29.	II	Prospective observational study. 365 patients. Results: Median time successful cannulation 3.2 min. 86% scene cannulation attempts successful. Conclusions and Recommendations: Paramedic training should ensure personnel can perform procedures quickly and that they are efficient in their use of on scene time.
Pons et al. ²³	1988	Prehospital venous access in an urban paramedic system—a prospective on-scene analysis. <i>J Trauma.</i> 1988;28:1460–1463.	II	Prospective observational study of timing of obtaining venous access. 125 patients. Results: Mean time to obtain access and sample blood was 2.2 ± 0.2. Blood draws add >60 s to venous access placement. First attempt success 90% and 100% of patients were able to get access. Statistical Methods: Descriptive data only. Conclusions/Recommendations: Patients should get access placed.
Gausche et al. ¹⁵	1998	Out-of-hospital intravenous access: unnecessary procedures and excessive cost. <i>Acad Emerg Med.</i> 1998;5:878–882.	II	Retrospective study of 452 consecutive patients admitted to an urban hospital. 84% received an IV (IV line or saline lock); 7% received fluid resuscitation in the field. 37% received “appropriate” treatment; 56% received an IV line when saline lock was indicated; 7% received saline lock when IV line needed. Conclusions/Recommendations: Paramedics and base hospital personnel often provide discordant over-treatment of patients by placement of an IV when a saline lock or no IV would suffice, resulting in unnecessary costs for EMS systems.
Sampalis et al. ²⁴	1997	Ineffectiveness of on-site intravenous lines: is prehospital time the culprit? <i>J Trauma.</i> 1997;43:608–615.	II	Observational study. 434 patients. Methods: 217 patients (IV group) compared with 217 patients (no IV) matched by Prehospital Index Score. Results: Mortality rates for the IV and no-IV groups were 23% and 6% ($p < 0.001$). Logistic regression analysis demonstrated on-site fluid replacement associated with significant increase in risk of mortality (adjusted OR = 2.3; 95% CI = 1.02–5.28; $p = 0.04$). Recommendations: Use of on-site IV fluid replacement is associated with an increase in mortality risk; association is exacerbated by increased prehospital time.
Smith et al. ²⁷	1985	Prehospital stabilization of critically injured patients: a failed concept. <i>J Trauma.</i> 1985;25:65–70.	III	Retrospective review of 52 patients divided into three groups. No SBP, SBP <70, SBP 70–100 mm Hg. In all groups, time to obtain access greater than transport time. No more than 450 mL of fluid received. No change in vital signs during transport with this intervention. Conclusion: Minimize field maneuvers, “scoop and run.”

TABLE 1. Should Vascular Access Be Obtained? (continued)

Author(s)	Year	Title	Class	Comments and Consensus
Seamon et al. ²⁵	2007	Prehospital procedures before emergency department thoracotomy: "scoop and run" saves lives. <i>J Trauma</i> . 2007;63:113–120.	III	Retrospective chart review. 180 consecutive penetrating trauma patients (2000–2005) who underwent ED thoracotomy. Patients divided into two groups by mode of transportation. Results: 88 patients arrived by emergency medical services (EMS), 92 by police or private vehicle (POV). Groups similar with respect to demographics. 8.0% EMS patients survived until hospital discharge vs. 17.4% of POV patients. Prehospital procedures performed in 88.6% EMS patients. Multivariate analyses identified prehospital procedures as the sole independent predictor of mortality. For each procedure, patients were 2.63 times more likely to die before hospital discharge (OR 0.38).
Slovic et al. ²⁶	1990	Success rates for initiation of intravenous therapy en route by prehospital care providers. <i>Am J Emerg Med</i> . 1990;8:305–307.	III	Retrospective evaluation of the ability to start venous access in the back of a rolling motor vehicle. 641 patients. Results: At least one IV line was started in 92% of trauma patients, regardless of blood pressure. In hypotensive patients, success rates for at least one IV was 95%. Average on scene time for all calls was 14.9 min (SD ± 10.1), whereas the average on scene time for hypotensive patients was 11.6 min (SD ± 6.2). Statistical Methods: Descriptive. Conclusions/Recommendations: Venous access can be secured with a high degree of success en route. Prompt transport of unstable patients should not be delayed solely to obtain IV access.
Spaite et al. ²⁸	1991	The impact of injury severity and prehospital procedures on scene time in victims of major trauma. <i>Ann Emerg Med</i> . 1991;20:1299–1305.	III	Prospective, observational study. Methods: Compared time on scene and evaluation for differences caused by severity of injury and procedures performed. 98 patients. Results: More severely injured patients had more procedures. Number of procedures did not increase scene time. Conclusions: In a highly developed trauma system, prehospital ALS procedures do not increase time to ED arrival.
Henderson et al. ¹⁷	1998	Unnecessary intravenous access in the emergency setting. <i>Prehosp Emerg Care</i> . 1998;2:312–316.	III	Retrospective chart review. 940 patients. Methods: Patients presenting to ED by EMS or private vehicle. 62% patients who had access placed by EMS did not have any benefit. 58% of ED placed lines went unused (in EMS patient) 52% of ED placed lines for private vehicle patients went unused. Conclusions: Many routinely placed lines are unnecessary and protocols should be studied to reduce this rate.
Cayten et al. ¹⁴	1993	Basic life support versus advanced life support for injured patients with an injury severity score of 10 or more. <i>J Trauma</i> . 1993;35:460–466.	III	Retrospective, multi-institutional study. 781 patients. Methods: Consecutive trauma patients with ISS ≥10. Subset analysis of 219 hypotensive patients. Results: 434 patients transported by ALS ambulance, 347 by BLS. No significant differences between ALS and BLS with regard to age or ISS. Observed survival in penetrating injury compared more favorably with MTOS predicted survival in BLS patients. Among hypotensive patients, observed survival also compared more favorably with MTOS predicted survival in BLS. Total prehospital times were not different between BLS and ALS. No benefit of ALS for trauma patients with total prehospital times of <35 min.
Hedges et al. ¹⁶	1988	Factors contributing to paramedic onscene time during evaluation and management of blunt trauma. <i>Am J Emerg Med</i> . 1988;6:443–448.	III	Retrospective cohort. 109 patients. Methods: Divided into high and low trauma score (TS) groups. Step-wise linear regression used to identify factors having effect on scene time. Results: Mean onscene time did not differ between high (>13) and low (≤13) TS groups. Higher number of procedures performed in the high TS patient groups. Patient groups with low TS showed no improvement in score with increasing on scene time. Conclusions: Paramedics tend to spend more time on scene when long transport times are initiated.
Consensus Working Group on Pre-hospital Fluids. ⁷	2001	Fluid resuscitation in pre-hospital trauma care: a consensus view. <i>J R Army Med Corps</i> . 2001;147:147–152.	III	Study design: Consensus group. Methods: None. Results: (1) Control of bleeding is paramount. (2) No access for superficial wounds. (3) Use mental status and radial pulse as initial triage means (start IV but hold fluids if present). (4) Start IV and provide 500 mL of colloid if no radial pulse or mental status not coherent. (5) Stop fluids if pulse and mental status return. (6) If no response, repeat 500 mL of colloid (hetastarch). Conclusions: Algorithm base on expert experience and literature from other military conflicts.

TABLE 2. If Access Is Obtained, Where and How Should It Be Placed?

Author(s)	Year	Title	Class	Comments and Consensus
Westfall et al. ⁴³	1994	Intravenous access in the critically ill trauma patient: a multicentered, prospective, randomized trial of saphenous cutdown and percutaneous femoral access. <i>Ann Emerg Med.</i> 1994;23:541-545.	I	Prospective, multicenter trial. Randomized, nonblinded, waiver of consent. 78 patients. Results: Percutaneous femoral vein cannulation faster than saphenous cut-down in patients with femoral pulse. Conclusions: Placement of an 8.5 F percutaneous femoral line is an acceptable alternative to saphenous vein cut-down in unstable trauma patients.
Guisto and Iserson. ³⁶	1990	The feasibility of 12-gauge intravenous catheter use in the prehospital setting. <i>J Emerg Med.</i> 1990;8:173-176.	II	Prospective observational. 38 patients. Methods: Patients evaluated over a 6-mo period, for access attempts. Results: There were 43 attempts at 12-gauge intravenous catheter placement in 38 patients. Overall success rate was 84% with a success-per-attempt rate of 74%. Conclusions/Recommendations: Paramedics can successfully use 12-gauge catheters.
Frascone et al. ³⁵	2007	Consecutive field trials using two different intraosseous devices. <i>Prehosp Emerg Care.</i> 2007;11:164-171.	II	Observational study. 178 patients. Results: Evaluation of intraosseous insertions (89 each of FAST 1TM and EZ-IO). Results: FAST 1TM—64 of 89 successful initial attempts (this group had more initial IV attempts as well). EZ-IO—78 of 89 successful initial attempts and were placed faster. Recommendations: If intraosseous access used, EZ-IO has higher success rate of insertion.
Benumof et al. ²⁹	1983	A large catheter sheath introducer with an increased side-port functional gauge. <i>Crit Care Med.</i> 1983;11:660-662.	III	Observational study. 17 patients. Methods: In vivo and in vitro experiments to judge flow rates of various catheters and introducers Results: High rates of infusion can be achieved with introducer catheters. Similarly, high rates of flow can be achieved via 14-gauge peripheral lines and through the side port of introducers as a 16-gauge line. Conclusions: Introducer side ports can function as volume lines even with indwelling pulmonary artery catheters (or slicks) in place.
Herron et al. ³⁷	1997	8.5 French peripheral intravenous access during air medical transport of the injured patient. <i>Air Med J.</i> 1997;16:7-10.	III	Retrospective review. 23 patients. Results: Evaluated injured patients who received a peripheral 8.5 Fr introducer during the study period. Initial small bore access achieved and this was exchanged over wire to larger catheter. Conclusions: Peripheral 8.5 Fr IV access via guide-wire exchange of an existing IV is a rapid and simple approach to large-bore IV access.

(b) An IV placed to “saline-lock” is equivalent in patency and function to a continuous infusion.

- C. Level III: (a) Intravenous fluid resuscitation should be withheld until active bleeding/hemorrhage is addressed. (b) Intravenous fluid administration in the prehospital setting (regardless of mechanism or transport time) should be titrated for palpable radial pulse using small boluses of fluid (250 mL) rather than fixed volumes or continuous administration.

If fluid is given, which type of fluid should be chosen? (Table 4).⁴⁴⁻⁵²

- A. Level I: (a) There is insufficient data to recommend one solution or type of fluid over other options in the prehospital setting. (b) Small volume boluses (250 mL) of 3% and 7.5% hypertonic saline (HTS) are equivalent, with respect to vascular expansion and hemodynamic changes, to large volume boluses (1 L) of standard solutions such as lactated Ringer’s (LR) or 0.9% normal saline (NS).
- B. Level II: There is insufficient data to support any recommendation at this level.
- C. Level III: There administration of blood in the prehospital setting is safe and feasible.

If fluid is given, how much and how fast should it be administered? (Table 5).^{10,39,42,50,51}

- A. Level I: No level one recommendation can be made. There is insufficient data to recommend specific rates or volumes of fluids to be administered in the prehospital setting.
- B. Level II: Fluids run at “keep vein open” rates are adequate for transporting injured patients.
- C. Level III: Rapid infusion systems and/or pressurized systems (to deliver fluids more rapidly) should not be used in the prehospital setting.

SCIENTIFIC FOUNDATION

General

On the heels of an ever expanding Prehospital Trauma Life Support curriculum, we have seen not only an expansion in the procedural skills set of EMS providers, but also a dramatic increase in the number of procedures performed. Although there is evidence to suggest potential benefit of prehospital procedures in rural blunt trauma patients with prolonged transportation time, there is little (if any) data to

TABLE 3. If Access Is Obtained, Should Fluid Be Administered in the Prehospital Setting?

Author(s)	Year	Title	Class	Comments and Consensus
Turner et al. ⁴²	2000	A randomised controlled trial of prehospital intravenous fluid replacement therapy in serious trauma. <i>Health Technol Assess.</i> 2000;4:1–57.	I	Prospective randomized controlled study. 1,309 patients. Methods: Randomization to (1) standard of care, (2) withholding fluids until arrival to hospital unless transport time was to be more than 1 h. Results: 699 patients randomized to standard fluids according to existing protocols and 610 patients were randomized to no fluids. Extremely poor compliance with only 31% of patients who were supposed to receive fluids actually received fluids and 80% of patients who were supposed to have fluids withheld actually received them. Mortality was similar between the two groups. Longer transport time in the group randomized to give standard fluids. Conclusions: Authors concluded that protocols recommending prehospital fluid administration do no harm.
Bickell et al. ¹⁰	1994	Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. <i>N Engl J Med.</i> 1994;331:1105–1109.	I	Prospective, randomized trial. 598 patients. Methods: Patient with penetrating torso injuries were randomized to delayed resuscitation (no fluids until OR) and standard of care fluid resuscitation for EMS. Results: Delayed resuscitation versus standard resuscitation, 70% versus 62% survival ($P = 0.04$). Shorter length of stay and higher nadir hemoglobin. Conclusions: In patients with penetrating torso injury, delayed resuscitation is recommended in the prehospital setting.
Boyle and Kuntz. ³⁰	1994	Saline locks in prehospital care. <i>Prehosp Disaster Med.</i> 1994;9:190–192.	II	Observational study. 110 patients. Methods: Evaluated the use of saline locks instead of traditional IV placement and continuous infusion. Results: All patients who required fluid resuscitation were appropriately given IVs. Those patients who required medications only appropriately received SL. Patency adequate. Conclusions: The use of a saline lock is an alternative to the use of traditional IV/infusion in the prehospital setting.
Carducci and Stein. ³¹	1994	Intravenous maintenance with a saline lock intermittent infusion device in the prehospital environment. <i>Prehosp Disaster Med.</i> 1994;9:67–70.	II	Prospective, nonblinded. 70 patients. Methods: Prospective evaluation to evaluate the effectiveness of saline lock to maintain access. Conclusions: Saline lock is an effective, rapid, reliable method of maintaining intravenous access during trauma transport. Continuous infusion is unnecessary.
Dula et al. ³³	2002	Use of prehospital fluids in hypotensive blunt trauma patients. <i>Prehosp Emerg Care.</i> 2002;6:417–420.	II	Retrospective. 150 patients. Methods: Matched pairs, case-control study of hypotensive patients (initial SBP <90 mm Hg). SBP higher in fluid group. No difference in outcome in patients who received ≥ 500 or <500 mL fluid. Time to ED was lower in no fluid group (45 vs. 54 min, $p = 0.02$). Conclusions: Control of airway, breathing, immobilization of fractures, and control of external bleeding should have priority over IV fluid administration in prehospital setting.
Eckstein et al. ³⁴	2000	Effect of prehospital advanced life support on outcomes of major trauma patients. <i>J Trauma.</i> 2000;48:643–648.	III	Retrospective. 496 patients. Methods: All major trauma patients transported by paramedics to a Level I Trauma Center. Results: Survival among patients who received IV fluids was not significantly greater than for those who did not receive fluids. Average on-scene times for patients who received fluids was not significantly longer than those who did not receive fluids. Conclusions: ALS procedures performed by paramedics on major trauma patients do not prolong scene time, but do not improve survival.
Holcomb. ³⁸	2003	Fluid resuscitation in modern combat casualty care: lessons learned from Somalia. <i>J Trauma.</i> 2003;54:S46–S51.	III	Expert opinion/consensus statement. Algorithm provided base on expert experience and literature from other military conflicts. (1) Superficial wounds do not require immediate IV access or fluid resuscitation, (2) if the soldier is coherent and has a palpable radial pulse, place a saline lock IV, (3) if incoherent or no radial pulse, obtain IV access and start 500 mL hetastarch, (4) repeat bolus if no response, saline lock IV if response noted, (5) in patients with suspected head injuries, fluids should be titrated for SBP >90 mm Hg.
Consensus Working Group on Pre-hospital Fluids. ⁷	2001	Fluid resuscitation in pre-hospital trauma care: a consensus view. <i>J R Army Med Corps.</i> 2001;147:147–152	III	Study design: Consensus group. Methods: None. Results: (1) Control of bleeding is paramount. (2) No access for superficial wounds. (3) Use mental status and radial pulse as initial triage means (start IV but hold fluids if present). (4) Start IV and provide 500 mL of colloid if no radial pulse or mental status not coherent. (5) Stop fluids if pulse and mental status return. (6) If no response, repeat 500 mL of colloid (hetastarch). Conclusions: Algorithm base on expert experience and literature from other military conflicts.

TABLE 3. If Access Is Obtained, Should Fluid Be Administered in the Prehospital Setting? (continued)

Author(s)	Year	Title	Class	Comments and Consensus
Revell et al. ⁴⁰	2002	Fluid resuscitation in prehospital trauma care: a consensus view. <i>Emerg Med J.</i> 2002;19:494–498.	III	Consensus Group statement. Recommendations: (1) Cannulation should take place en route where possible, (2) only two attempts at cannulation should be made, (3) transfer should not be delayed by attempts to obtain intravenous access, (4) entrapped patients require cannulation at the scene, (5) normal saline is recommended as a suitable fluid for administration to trauma patients, (6) boluses of 250-mL fluid may be titrated against the presence or absence of a radial pulse (caveats; penetrating torso injury, head injury, infants).
Talving et al. ⁴¹	2005	Prehospital management and fluid resuscitation in hypotensive trauma patients admitted to Karolinska University Hospital in Stockholm. <i>Prehosp Disaster Med.</i> 2005;20:228–234.	III	Retrospective. 102 patients. Methods: A retrospective, descriptive study on consecutive, hypotensive trauma patients. Results: 75% blunt. On-scene time 19 min. Fluid therapy initiated at the scene in 73%, regardless of mechanism or injury severity. Conclusions: Time interval at the scene of injury exceeded PHTLS guidelines. Majority of hypotensive patients were fluid-resuscitated on-scene regardless of the mechanism or injury severity.
Dalton. ³²	1995	Prehospital intravenous fluid replacement in trauma: an outmoded concept? <i>J R Soc Med.</i> 1995;88:213P–216P.	III	Observational study. 235 patients. Methods: Injured patients admitted to trauma center (72% penetrating) who received prehospital IV placement. Results: Mean infusion time 17 min. 92% patients received volumes <1,000 mL, 80% received less than <600 mL. Hypotensive patients had mean infusion time of 16 min, receiving mean 573 mL. Conclusions: Given the uncertain benefits and potential complications, intravenous cannulation and fluid replacement may not be appropriate where expected prehospital time is likely to be less than 30 min.
Smith et al. ²⁷	1985	Prehospital stabilization of critically injured patients: a failed concept. <i>J Trauma.</i> 1985;25:65–70.	III	Retrospective review of 52 patients divided into three groups. No SBP, SBP <70, SBP 70–100 mm Hg. In all groups, time to obtain access greater than transport time. No more than 450 mL of fluid received. No change in vital signs during transport with this intervention. Conclusion: Minimize field maneuvers, “scoop and run.”
Kaweski et al. ³⁹	1990	The effect of prehospital fluids on survival in trauma patients. <i>J Trauma.</i> 1990;30:1215–1218.	III	Retrospective. 6,855 patients. Methods: Evaluation of a cohort of patients from a trauma registry. Results: 56% of patients received fluids, penetrating got fluid more often. Longer transport time in patients receiving IV and fluids. No difference in receiving fluids and mortality by ISS, TRISS, or hypotension on arrival. Conclusions: Prehospital fluids do not affect outcome but initiating fluids does delay transport.

justify these procedures in patients with penetrating injuries or those with short transport times (less than 30 minutes).^{10,27,53} Advanced Trauma Life Support guidelines for administering (up to 2 L) crystalloid to hypotensive trauma patients in the emergency room setting have been adopted into the prehospital environment without clear benefit. In addition, recent data has shown that the performance of prehospital procedures in urban and penetrating settings have a negative impact on survival.²⁵ In light of the expanding data calling into question the “stay and play” method of prehospital care, the EAST PMG committee conducted a review of the currently available literature on prehospital vascular access and fluid resuscitation.

Should Vascular Access Be Obtained in the Prehospital Setting?

The first issue examined was whether or not vascular access should even be attempted in the prehospital setting. Several investigators have evaluated the success rate for prehospital venous access and noted scene placement times of these lines range from a mean of 2.2 minutes to 6.3 minutes.^{18–20,23,27} “Time to placement” of these lines was similar in patients

with normal blood pressure and primary injuries confined to the extremities.^{13,15,22} However, when patients were hypotensive or had primarily torso injuries, placement at the scene tended to be longer than that of en route intravenous line placement.^{21,26,27} Jones et al.¹⁹ noted a 91% success rate at the scene and a 94% success rate en route. Additionally, Slovis et al.²⁶ demonstrated an en route success rate for intravenous line placement of 92%, regardless of hemodynamic status. Delaying transport to place venous access also seems to be associated with increased overall time to hospital, in some cases exceeding that of the actual transport itself.^{24,25,27} Should the decision be made to obtain scene access, drawing blood for laboratory samples (e.g., type and screen, etc) adds unnecessary time and should not be performed.²³

Whether the committee felt it was able to recommend placement of access in the prehospital setting was predicated on demonstrating any benefit from access placement. Although several studies evaluating prehospital venous access have recommended their placement (simply because they could technically be placed), none of the investigators was

TABLE 4. If Fluid Is Administered, Which Solution Should Be Given?

Author(s)	Year	Title	Class	Comments and Consensus
Vassar et al. ⁵¹	1991	7.5% sodium chloride/dextran for resuscitation of trauma patients undergoing helicopter transport. <i>Arch Surg.</i> 1991;126:1065–1072.	I	Randomized, double blind multicenter trial. 166 patients. Methods: Waiver of consent obtained. Study compared 250 mL boluses of LR and 7.5% + 6% Dextran. Remainder of resuscitation with standard solutions. Results: No significant difference in initial SBP, mortality, use of blood products overall. In patients with GCS score <8 hypertonic saline groups had had higher survival. Mean change in SBP was higher in hypertonic groups. Conclusions: Prehospital use of hypertonic saline is feasible in air medical transported patients and is associated with improved survival in those patients with head injury.
Vassar et al. ⁵⁰	1993	A multicenter trial for resuscitation of injured patients with 7.5% sodium chloride. The effect of added dextran 70. The Multicenter Group for the Study of Hypertonic Saline in Trauma Patients. <i>Arch Surg.</i> 1993; 128:1003–1011.	I	Randomized, double blind multicenter trial. 194 patients. Methods: Waiver of consent obtained. Critically injured patients transported by air medical ambulance. Compared 250 mL bolus of LR, 7.5% sodium chloride, 7.5% + 6% Dextran, or 7.5% + 12% Dextran. Remainder of resuscitation with standard solutions. Results: Hypertonic saline associated with improved admission SBP compared with LR. No differences noted with the addition of Dextran. Brain injury patients seem to benefit the most with 7.5% saline. Conclusions: Use of hypertonic saline, with or without Dextran, reduces mortality in trauma patients.
Vassar et al. ⁵²	1993	Prehospital resuscitation of hypotensive trauma patients with 7.5% NaCl versus 7.5% NaCl with added dextran: a controlled trial. <i>J Trauma.</i> 1993;34:622–632.	I	Randomized, provider blinded trial. 258 patients. Methods: Single-center trial of all patient transported by ground ambulance with SBP <90 en route. Received 0.9% sodium chloride, 7.5% sodium chloride, and 7.5% with Dextran. Results: ISS higher in 7.5% groups. 7.5% group had more unexpected survivors than 0.9% group but the overall mortality was not significantly different. No benefit to dextran. Conclusions: Infusion of small bolus of 7.5% saline may improve survival in severely injured patients.
Maningas et al. ⁴⁷	1989	Hypertonic saline-dextran solutions for the prehospital management of traumatic hypotension. <i>Am J Surg.</i> 1989;157:528–533.	I	Randomized, double-blinded controlled trial. 48 patients. Methods: All patients with penetrating injury and a SBP <90. Randomized in the field to 7.5% sodium chloride + 6% dextran or crystalloid (plasmalyte) based on an alternate-day protocol. Results: Trends toward improved SBP response and less blood transfusion with the study group; yet, neither reached statistical significance. Conclusions: It is feasible to give low volume solution in a prehospital setting. There were trends toward improvements in clinically significant outcomes in patients receiving 7.5% sodium chloride.
Mattox et al. ⁴⁸	1991	Prehospital hypertonic saline/dextran infusion for post-traumatic hypotension. The U.S.A. Multicenter Trial. <i>Ann Surg.</i> 1991;213:482–491.	I	Multicenter, randomized double-blinded trial. 359 patients. Methods: Waiver of consent, intention to treat trial. Randomized in the field to 7.5% sodium chloride + 6% dextran or crystalloid. Results: Nonstatistically significant trend to higher overall survival in treatment arm. However, significantly better survival was observed in patients requiring operative intervention ($p = 0.02$) and in those with penetrating injuries ($p = 0.01$). Conclusions: Hypertonic saline as safe and effective as standard resuscitation. This solution may improve survival in patients who require early operative intervention or in those who sustain penetrating injuries.
Cooper et al. ⁴⁵	2004	Prehospital hypertonic saline resuscitation of patients with hypotension and severe traumatic brain injury: a randomized controlled trial. <i>JAMA.</i> 2004;291:1350–1357.	I	Prospective, randomized, double blind trial. 262 patients. Methods: Community waiver of consent. Evaluated patients with either head injury or hypotension. Randomized to 7.5% sodium chloride or Ringer's lactate. Results: No difference in overall mortality or 6 mo neurological outcome. Lower mean ICP on admission in 7.5% group. Conclusions: Hypertonic saline is not associated with improved neurological outcomes at 6-mo.
Holcroft et al. ⁴⁶	1987	3% NaCl and 7.5% NaCl/dextran 70 in the resuscitation of severely injured patients. <i>Ann Surg.</i> 1987;206:279–288.	I	Randomized, provider blinded trial. 258 patients. Methods: Single-center trial of all patient transported by ground ambulance with SBP <90 en route. Received lactated Ringer's or 7.5% sodium chloride with dextran. Results: Overall survival was 40% in hypertonic group versus 30% in the LR group (not significant). The LR arm required more fluids to maintain SBP en route and in ED. 7.5% group had a greater increase in SBP from baseline (49 mm Hg versus 19 mm Hg, $p < 0.005$). Conclusions: Prehospital use of 7.5% sodium chloride with dextran improves SBP more than LR and may improve survival.
Barkana et al. ⁴⁴	1999	Prehospital blood transfusion in prolonged evacuation. <i>J Trauma.</i> 1999;46:176–180.	III	Retrospective chart review. 40 patients. Methods: Evaluated all cases of trauma patients in who received prehospital blood transfusions during a 30-mo period. Results: 40 patients received 60 units of blood. Mean prehospital crystalloid 4.4 L. Over the 30-mo period, less than 4% of blood units that were "on standby" for prehospital situations were actually used (90% used in hospital). Conclusions: Prehospital blood transfusion is feasible and safe. However, a large amount of wasted blood occurs using prehospital blood.
Sumida et al. ⁴⁹	2000	Prehospital blood transfusion versus crystalloid alone in the air medical transport of trauma patients. <i>Air Med J.</i> 2000;19:140–143.	III	Retrospective chart review. 48 patients. Methods: One-year review of air medical transported patients. Control group made up of those patients receiving no blood but >2.0 L crystalloid en route. Study group made up of patients who received in-flight transfusions. Results: 31 patients received crystalloid only; 17 received blood and crystalloid. No demographic differences between the groups. However, the crystalloid only group had higher initial blood pressure and heart rate and the blood group had longer transport times. In addition, the blood group had lower pH and HCO ₃ on admission. No difference in mortality. Conclusions: The impact of blood products in outcomes could not be assessed because of significant differences in transport times.

TABLE 5. If Fluid Is Administered, How Much Should Be Given and How Fast Should It Be Infused?

Author(s)	Year	Title	Class	Comments and Consensus
Vassar et al. ⁵¹	1991	7.5% sodium chloride/dextran for resuscitation of trauma patients undergoing helicopter transport. <i>Arch Surg.</i> 1991;126:1065–1072.	I	Randomized, double blind multicenter trial. 166 patients. Methods: Waiver of consent obtained. Study compared 250 mL boluses of LR and 7.5% + 6% Dextran. Remainder of resuscitation with standard solutions. Results: No significant difference in initial SBP, mortality, use of blood products overall. In patients with GCS score <8 hypertonic saline groups had had higher survival. Mean change in SBP was higher in hypertonic groups. Conclusions: Prehospital use of hypertonic saline is feasible in air medical transported patients and is associated with improved survival in those patients with head injury.
Vassar et al. ⁵⁰	1993	A multicenter trial for resuscitation of injured patients with 7.5% sodium chloride. The effect of added dextran 70. The Multicenter Group for the Study of Hypertonic Saline in Trauma Patients. <i>Arch Surg.</i> 1993;128:1003–1011.	I	Randomized, double blind multicenter trial. 194 patients. Methods: Waiver of consent obtained. Critically injured patients transported by air medical ambulance. Compared 250 mL bolus of LR, 7.5% sodium chloride, 7.5% + 6% Dextran, or 7.5% + 12% Dextran. Remainder of resuscitation with standard solutions. Results: Hypertonic saline associated with improved admission SBP compared to LR. No differences noted with the addition of Dextran. Brain injury patients seem to benefit the most with 7.5% saline. Conclusions: Use of hypertonic saline, with or without Dextran, reduces mortality in trauma patients.
Turner et al. ⁴²	2000	A randomised controlled trial of prehospital intravenous fluid replacement therapy in serious trauma. <i>Health Technol Assess.</i> 2000;4:1–57.	I	Prospective randomized controlled study. 1,309 patients. Methods: Randomization to (1) standard of care, (2) withholding fluids until arrival to hospital unless transport time was to be more than 1 h. Results: 699 patients randomized to standard fluids according to existing protocols and 610 patients were randomized to no fluids. Extremely poor compliance with only 31% of patients who were supposed to receive fluids actually received fluids and 80% of patients who were supposed to have fluids withheld actually received them. Mortality was similar between the two groups. Longer transport time in the group randomized to give standard fluids. Conclusions: Authors concluded that protocols recommending prehospital fluid administration do no harm.
Bickell et al. ¹⁰	1994	Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. <i>N Engl J Med.</i> 1994;331:1105–1109.	I	Prospective, randomized trial. 598 patients. Methods: Patient with penetrating torso injuries were randomized to delayed resuscitation (no fluids until OR) and standard of care fluid resuscitation for EMS. Results: Delayed resuscitation versus standard resuscitation, 70% versus 62% survival ($p = 0.04$). Shorter length of stay and higher nadir hemoglobin. Conclusions: In patients with penetrating torso injury, delayed resuscitation is recommended in the prehospital setting.
Kaweski et al. ³⁹	1990	The effect of prehospital fluids on survival in trauma patients. <i>J Trauma.</i> 1990;30:1215–1218.	III	Retrospective. 6,855 patients. Methods: Evaluation of a cohort of patients from a trauma registry. Results: 56% of patients received fluids, penetrating got fluid more often. Longer transport time in patients receiving IV and fluids. No difference in receiving fluids and mortality by ISS, TRISS, or hypotension on arrival. Conclusions: Prehospital fluids do not affect outcome but initiating fluids does delay transport.

able to demonstrate any benefit from their placement.^{18,20,23} To the contrary, scene access placement has been associated with increased risk of death.^{14,24,27} Seamon et al.²⁵ evaluated 180 patients who underwent ED thoracotomy over a 6-year period. The authors stratified according to method of transport (EMS or private vehicle/police [POV]). They found that mortality in patients arriving by EMS was double that of patients transported by POV (17% vs. 8.0%). More impor-

tantly, each prehospital procedure performed reduced the odds of survival by 62% (OR, 0.38; 95% CI, 0.18–0.79; $p = 0.009$). Sampalis et al.²⁵ examined the association between on-site venous access and fluid replacement and mortality in critically injured patients. The authors found that on-site access was associated with an increase in mortality (23% IV group vs. 6% no IV group) and that this association was stronger as time to hospital arrival increased (OR, 8.4; CI,

1.27–54.69; $p = 0.03$ for time to arrival >60 minutes). These outcomes (as well as time to placement of access and scene times) do not seem to differ whether the patient is transported by basic life support (BLS) or advanced life support ambulance crews.

In light of increasing experience in management of military injuries (both penetrating and blunt), a UK consensus group was recently convened to reconcile differences in practice with respect to prehospital fluid resuscitation.⁷ The expert panel formulated an algorithm to assist in the early resuscitation of critically injured patients that consisted of the following: (1) access should not be obtained for superficial wounds, (2) if the patient's mental status is appropriate and a radial pulse is present, prehospital personnel may place venous access but fluids should be held, (3) venous access should be obtained and fluids initiated if no radial pulse or mental status is incoherent, (4) a 500 mL bolus of hetastarch should be given if no radial pulse or mental status is incoherent, (5) fluids should be discontinued if pulse and mental status return, and (6) if no response, repeat 500 mL of colloid (hetastarch).⁷

If Access Is Obtained, Where Should It Be Placed?

In asking where to place venous access, it is important not only to determine where anatomically should this access be obtained, but also what approach should be used (percutaneous, cut-down) and what size catheter should be placed. As well, it is critical to evaluate whether there are alternatives to traditional intravenous catheter access (intraosseous lines). No clinical trials were found to suggest superiority of one anatomic location for venous access versus another. Peripheral access has been the standard location for placement in the prehospital setting; most commonly being placed in the upper extremity in the forearm and antecubital area. An animal study by Rosa et al. examined the efficacy of intravenous access site on delivery of a bolus injection to the heart during shock and resuscitation.⁵⁴ The authors noted that femoral access significantly prolongs bolus transit time when compared with central or brachial access regardless of intravascular status (euvolemic, hypovolemic, or aggressively resuscitated). Brachial access was advocated as the preferred route for bolus injection delivery in the emergency setting as it provides expedient bolus delivery equal to central access and is superior to femoral access.

A few authors have evaluated the time-to-placement and efficiency of traditional peripheral access, cut-down placement, and percutaneous (Seldinger) approaches in the prehospital setting.^{29,36,43} Westfall et al.⁴³ conducted a prospective, randomized, multicentered trial to compare the speed of IV access and the rate of infusion for saphenous vein cut-down and percutaneous femoral catheterization. Seventy-eight trauma patients were randomized to either saphenous cut-down or percutaneous femoral line placement, followed by passive infusion of 1.0 L of crystalloid. Mean procedure time for the cut-down group was 5.6 minutes \pm 2.6 minutes compared with 3.2 minutes \pm 1.2 minutes for the femoral line group ($p < 0.001$). The mean infusion time for the cut-down group was 6.6 minutes \pm 4.3 minutes compared with 4.6

minutes \pm 2.5 minutes for the femoral line group ($p = 0.03$). As the rate of flow increases with larger diameter and shorter length of the catheter (Poiseuille's law), several investigators have examined the impact of placing larger bore lines in the prehospital setting. Benumof et al.²⁹ measured the infusion pressure-flow characteristics of the side ports of 8.0 and 8.5 Fr introducer catheters against equivalent pressure-flow relationship through standard 22- to 14-gauge peripheral venous catheters. Although the 8.5 Fr introducers were superior to the 8.0 Fr catheters, none was superior to 14- and 16-gauge standard peripherally placed catheters. Other authors evaluated the ability to place extremely large bore peripheral lines (10- and 12-gauge catheters) and to rapidly exchange smaller peripheral lines out to large introducer catheters (over a wire).^{36,37} However, none demonstrated superiority to standard large bore (14–18 gauge) peripheral lines.

Over the last several years, intraosseous lines have been met with increasing enthusiasm and favor in the prehospital resuscitation of trauma patients. However, this enthusiasm is based on increasing provider "experiences" and not on any clinical trials. These studies are, for the most part, are limited to "how I do it" or device comparisons, using historical flow rates of intravenous catheters as reference values.⁵⁵ Others are pharmacokinetic drug studies or trials conducted in nontrauma patients. Frascone et al.³⁵ studied provider performance for obtaining intraosseous access with two FDA-approved intraosseous devices in two sequential field trials. They evaluated 178 insertions in adult trauma patients, with success rates of 72% to 87%. Time to insertion was similar to that of historical data on intravenous access. The site most often used for adult intraosseous access is the proximal tibia (medial and inferior to the tibial tuberosity) and the sternum and humeral head.⁵⁵ Although the pharmacokinetic delivery of drugs seems equal to that of the intravenous route, infusion volume rates are only that of a 21-gauge catheter in the absence of a high-pressure infusion system.^{55,56}

If Access Is Obtained, Should Fluid Be Administered in the Prehospital Setting?

Once access is established, another controversy arises in whether or not to initiate fluid therapy.³⁴ Although many providers see the subsequent administration of fluids to be standard of care, the literature would not support this approach. At a minimum, prehospital fluid administration does not seem to improve outcomes in either penetrating or blunt trauma patients.^{27,32,39} In a study of 235 trauma patients (blunt and penetrating), Dalton³² evaluated the benefit of prehospital venous access and fluid administration. The authors noted that 80% patients receive less than 600 mL of fluid in the prehospital setting, regardless of mechanism, scene entrapment, or hypotension en route. They were unable to identify benefit from such therapy and recommended withholding fluid administration. Kaweski et al.³⁹ conducted a retrospective study of almost 7,000 trauma patients and noted that mortality rates were similar in those patients who received fluids and those who did not (23% vs. 22%; $p = NS$). Comparison of groups with similar, injury severity,

probability of survival, and hypotension on arrival also failed to show an influence of fluid administration on survival. Other authors have also demonstrated that prehospital fluid therapy confers no survival benefit and delays transport of critically injured patients.^{33,41}

Although numerous animal studies have demonstrated an increased rate of hemorrhage and an increased mortality with “prehospital” fluid resuscitation, clinical extrapolation of this concept has, for the most part, been confined to a single prospective randomized controlled trial by Bickell et al.¹⁰ in the early 1990s. Hypotensive patients with penetrating torso injuries were randomized in the field to receive either standard intravenous fluid resuscitation or no fluids, and the regimen was continued until the patient reached the operating theater. Despite the lack of prehospital fluid resuscitation, the mean blood pressure between the groups was similar while intraoperative blood loss was significantly less in the delayed resuscitation group. Additionally, the length of stay was shorter and the mortality rate lower in the delayed resuscitation group. Another randomized controlled trial by Turner et al.⁴² in the UK involved 1,309 patients, half of which were randomized to standard fluid protocols or no prehospital fluids. Unfortunately, not much can be derived from this study to make any recommendations, as there was extremely poor protocol compliance. Only 31% of standard protocol patients (who were supposed to receive fluids) received prehospital fluids and only 80% of the “no fluid” patients had fluids withheld. There was no significant difference in mortality between the groups. Despite the studies significant limitations, the authors concluded that prehospital fluid resuscitation does no harm.

Several groups and authors have developed consensus statements aimed directly at the question of whether to place venous access in the prehospital setting and whether to administer fluids before hospital arrival (and definitive hemorrhage control).^{7,37,55}

In summary, the consensus recommendations are quite similar: (1) patients with only superficial wounds (even in combat settings) do not require immediate intravenous access or fluid resuscitation, (2) if the patient is coherent and has a palpable radial pulse, place the venous access to “saline lock,” (3) if incoherent or no radial pulse, obtain venous access and start 500 mL hetastarch, (4) repeat bolus if no response, saline lock IV if response noted, (5) in patients with suspected head injuries, fluids should be titrated for SBP >90 mm Hg. These recommendations reflect that of the groups’ evaluations of blunt and penetrating patients in both military and civilian settings.⁴⁰

With respect to maintaining venous access patency, several authors have evaluated whether continuous intravenous infusion is necessary.^{30,31} Boyle and Kuntz³⁰ evaluated 100 patients requiring intravenous access placement. The use of a saline lock was found to be a cost-effective means of maintaining patency of intravenous lines during transport. Carducci and Stein³¹ demonstrated that a saline lock device was as effective at maintaining prehospital access as were traditional continuous infusions. Additionally, paramedics found that the saline lock devices were easier to use, less time-consuming to initiate, and facilitated patient transportation.

If Fluid Is Administered, Which Solution Should Be Given?

Although “standard” fluid on prehospital vehicles is typically normal saline (0.9% sodium chloride NS) or lactated Ringer’s (LR) solution, neither has been well studied in the prehospital resuscitation of trauma and neither has been shown to be superior to other available solutions. However, their cost and provider familiarity with these solutions has likely been responsible for their “standard solution” status. Studies comparing various iso-oncotic solutions (hetastarch, dextran) and hypertonic saline (3% or 7.5% sodium chloride) to NS or LR have shown mixed results.^{46,47,50,52,57–61}

Cooper et al.⁴⁵ recently compared LR and 7.5% sodium chloride in a randomized trial of trauma patients with either brain injury or hypotension. The authors found no significant difference between the groups with respect to favorable neurologic outcomes at 6 months. Survival to discharge and 6-month survival were higher in the 7.5% sodium chloride group (55% vs. 50% and 55% vs. 47%, respectively), but this did not reach statistical significance (study powered to detect difference in neurologic outcomes). In a trial of patients undergoing air medical transport after injury, Vassar et al.^{51,52} compared 7.5% sodium chloride (with and without dextran) with LR solution in a randomized fashion. The authors found that patients who received the hypertonic saline solutions had less fluid requirements in the prehospital setting and arrived to the trauma center with a higher systolic blood pressure. Although overall survival was not different between the hypertonic saline and LR groups, patients with severe head injury who received hypertonic saline had a higher survival than that observed with the LR groups (32% vs. 16%; $p = 0.044$). The same group evaluated 0.9% sodium chloride (NS) with several hypertonic saline solutions in hypotensive patients transported by ground ambulance.⁵² In this randomized, controlled and prehospital blinded trial, the authors found no difference in survival between patients who received 0.9% or 7.5% sodium chloride. However, the groups were poorly matched with significantly higher injury severity scores and predicted mortality in the 7.5% groups compared with the 0.9% group (both $p < 0.05$). In addition, the study was severely underpowered with an estimated sample size calculation required of almost 700 patients (study had 258 patients).

Numerous investigators have examined the different hypertonic solutions available, including 3% and 7.5% sodium chloride and solutions with and without added colloids (6% and 12% dextran).^{45–47,50–52,57–59,62–64} With respect to 3% versus 7.5% sodium chloride, there are few, if any, animal studies comparing the solutions and no clinical trials of prehospital resuscitation of trauma. In a porcine model of uncontrolled hemorrhage, Watters et al.⁶⁴ recently evaluated the effect of 3% versus 7.5% sodium chloride. The authors noted that a single bolus of 3% solution produced an adequate and sustained rise in blood pressure and tissue oxygenation, whereas 7.5% sodium chloride failed to produce a sustained improvement in these variables. Also, the 7.5% sodium chloride solution resulted in a significant dilutional anemia and relative hypofibrinogenemia. Prehospital clinical trials have utilized

7.5% sodium chloride (with and without dextran) almost exclusively, thus preventing a critical comparison of and evidenced-based recommendation by the committee for one over the other.^{46–48,50–52,58} As to the benefit of adding colloid to hypertonic saline, solutions of 7.5% sodium chloride with 6% or 15% dextran added do not seem superior to those without dextran.^{50–52} Vassar et al.⁵⁰ noted that the use of 7.5% sodium chloride was superior to LR in reducing the amount of prehospital fluid requirements and improving admission/emergency department blood pressure, with no additional benefit seen with the dextran solutions. Glasgow outcome scores were higher in the 7.5% group compared with lactated Ringer's, but no further effect was observed with the dextran solutions.

Hydroxylethylstarch (HES) is a balanced electrolyte solution that is similar in ionic composition to human plasma.^{65,66} Although it is contraindicated in patients with bleeding disorders, HES is used and advocated by the military for prehospital, low-volume boluses of injured soldier's in hemorrhagic shock.^{38,40} In fact, several near-fatal hemorrhage models have demonstrated that HES is associated with hemorrhage volume and significantly lower mortality when compared with LR.^{47,61–63} Compared with normal saline and lactated Ringer's solution, HES is associated with significant decreases in release of proinflammatory cytokines.^{19,62–64} When compared with LR, HES solutions in animals with trauma-hemorrhagic shock has been shown to induce less inflammatory cytokines and improved immune function.

Clinical trials evaluating blood and blood substitutes in the prehospital setting are also scarce and prevented the committee from making any significant recommendation on the subject. Barkana et al.⁴⁴ evaluated 40 patients who had received prehospital blood transfusions. Prolonged extrication and delayed transport were the primary indications for transfusion and the mean volume of crystalloid infused by hospital arrival was almost 4.5 L. During the study period, only 4% of blood reserved for prehospital patients was actually used (vs. 90% during that period for in-hospital use). Although the authors found it safe and feasible to transfuse blood in the prehospital setting, there was a tremendous amount of product wastage. Sumida et al.⁴⁹ performed a chart review of patients receiving prehospital resuscitation with blood products and found no difference in mortality. The authors noted that the crystalloid only group, however, had shorter transport times and more "normal" vital signs in the field. As to blood substitutes, data regarding use in trauma patients are extremely limited while that for prehospital resuscitation are essentially nonexistent.^{67,68} Gould et al.⁶⁵ conducted a prospective, randomized, open label trial of a human polymerized hemoglobin substitute. Forty-four in-hospital trauma patients were randomized to receive red cells or up to six units of the blood substitute as their initial blood replacement after trauma and during emergent operations. The total number of allogeneic red cell transfusions for the control and experimental groups was not significantly different through study day 3 (11.3 ± 4.1 units vs. 7.8 ± 4.2 units; $p = 0.06$).

If Fluid Is Administered, How Much Should Be Given and How Fast Should It Be Infused?

Even more poorly understood and more under-studied than the above controversies is that of "how much" and "how fast." Basic trauma guidelines recommend an initial rapid infusion of fluid (1.0–2.0 L) in hypotensive trauma patients as a diagnostic procedure to aid treatment decisions. Although this has been interpreted (or misinterpreted) to mean that patients in the prehospital setting should receive 2.0 L of crystalloid (often regardless of hemodynamic status), this practice is not supported by the literature. Currently, there are no clinical trials to support a recommendation for rate or volume of fluid administration to trauma patients in the prehospital setting. Animal studies have been performed that suggest higher volumes and more rapid infusions of crystalloid in the prehospital setting result in higher mortality.^{63,64,66,69–71} Using a murine model of uncontrolled hemorrhage, Krausz et al.⁷² compared bolus and continuous infusion administration of lactated Ringer's and hypertonic saline, combined with splenectomy. Continuous infusion of LR solution resulted in significantly less bleeding than bolus infusion and improved survival time, whereas there was no difference in continuous or bolus infusion of hypertonic saline. Bickell et al.¹⁰ demonstrated that patients with penetrating torso injury who received small volumes of resuscitation (<250 mL) in the prehospital setting had significantly higher survival to hospital discharge than those who received standard prehospital resuscitation (750–1,000 mL in less than 30 minutes). Vassar et al.^{50–52} noted that hypertonic saline boluses (on the order of 250 mL volumes) seem to be associated with improved survival.

In-patient clinical trials suggest that "lowering expectations" of resuscitation end-points may improve survival and decrease hemorrhage volume. Dutton et al.⁹ evaluated the concept of "hypotensive resuscitation" and noted that titration of initial fluid therapy to a lower than normal systolic blood pressure (SBP) (≥ 70 mm Hg instead of >100 mm Hg) during active hemorrhage did not affect mortality. Several years earlier, this same group of investigators evaluated the delivery of crystalloid and blood products through a rapid infusion system to critically injured patients in hemorrhagic shock.⁷³ Contrary to beliefs and biases at the time, the investigators noted an almost fivefold increase in mortality in patients resuscitated with a rapid infusion system. This higher mortality remained even after matching for age, injury severity scores, and Glasgow coma scale (53% vs. 61%; $p < 0.001$). Even more surprising was that the mortality difference was greatest not in penetrating patients but in the blunt trauma population (48.8% vs. 63.0%; $p < 0.001$).

CONCLUSION

Despite the widely held belief that prehospital venous access placement and fluid resuscitation is standard of care, there is little data to support this practice. In fact, an increasing amount of data suggests that it may be quite harmful to a significant number of critically injured patients. The EAST PMG committee has found that placement of venous access at the scene delays transport and placement of access en route

should be considered. In those patients in whom intravenous access has failed, intraosseous may be attempted if equipment and trained personnel are available. There is insufficient data to suggest that blunt or penetrating trauma patients benefit from prehospital fluid resuscitation. In patients with penetrating injuries and short transport times (less than 30 minutes), fluids should be withheld in the prehospital setting in patients who are alert or have a palpable radial pulse. Fluids (in the form of small boluses, i.e., 250 mL) should be given to return the patient to a coherent mental status or palpable radial pulse. In the setting of traumatic brain injury, however, fluids should be titrated to maintain systolic blood pressure greater than 90 mm Hg (or mean pressure greater than 60 mm Hg). Hypertonic saline boluses of 250 mL seem equivalent in efficacy to 1,000 mL boluses of standard solutions (lactated Ringer's, 0.9% sodium chloride). There is insufficient evidence to show that injured patients with short transport times benefit from prehospital blood transfusions. Finally, rapid infusion systems and or pressurized devices (to deliver fluids more rapidly) should not be used in the prehospital setting.

REFERENCES

- Salomone JP, Ustin JS, McSwain NE Jr, Feliciano DV. Opinions of trauma practitioners regarding prehospital interventions for critically injured patients. *J Trauma*. 2005;58:509–515.
- Bochicchio GV, Ilahi O, Joshi M, Bochicchio K, Scalea TM. Endotracheal intubation in the field does not improve outcome in trauma patients who present without an acutely lethal traumatic brain injury. *J Trauma*. 2003;54:307–311.
- Wang HE, Peitzman AB, Cassidy LD, Adelson PD, Yealy DM. Out-of-hospital endotracheal intubation and outcome after traumatic brain injury. *Ann Emerg Med*. 2004;44:439–450.
- Cornwell EE III, Belzberg H, Hennigan K, et al. Emergency medical services (EMS) vs non-EMS transport of critically injured patients: a prospective evaluation. *Arch Surg*. 2000;135:315–319.
- Demetriades D, Chan L, Cornwell E, et al. Paramedic vs. private transportation of trauma patients. Effect on outcome. *Arch Surg*. 1996;131:133–138.
- Liberman M, Mulder D, Lavoie A, Denis R, Sampalis JS; Multicenter Canadian study of prehospital trauma care. *Ann Surg*. 2003;237:153–160.
- Fluid resuscitation in pre-hospital trauma care: a consensus view. *J R Army Med Corps*. 2001;147:147–152.
- Dretzke J, Sandercock J, Bayliss S, Burls A. Clinical effectiveness and cost-effectiveness of prehospital intravenous fluids in trauma patients. *Health Technol Assess*. 2004;8:iii1–iii103.
- Dutton RP, Mackenzie CF, Scalea TM. Hypotensive resuscitation during active hemorrhage: impact on in-hospital mortality. *J Trauma*. 2002;52:1141–1146.
- Bickell WH, Wall MJ Jr, Pepe PE, et al. Immediate versus delayed fluid resuscitation for hypotensive patients with penetrating torso injuries. *N Engl J Med*. 1994;331:1105–1109.
- Oxman AD, Cook DJ, Guyatt GH. Users' guides to the medical literature. VI. How to use an overview. Evidence-Based Medicine Working Group. *JAMA*. 1994;272:1367–1371.
- Oxman AD, Schunemann HJ, Fretheim A. Improving the use of research evidence in guideline development. XIV. Reporting guidelines. *Health Res Policy Syst*. 2006;4:26.
- Barrett B, Guly HR. How long does it take to perform procedures on scene? *Prehospital Inmed Care*. 2000;4:25–29.
- Cayten CG, Murphy JG, Stahl WM. Basic life support versus advanced life support for injured patients with an injury severity score of 10 or more. *J Trauma*. 1993;35:460–466.
- Gausche M, Tadeo RE, Zane MC, Lewis RJ. Out-of-hospital intravenous access: unnecessary procedures and excessive cost. *Acad Emerg Med*. 1998;5:878–882.
- Hedges JR, Feero S, Moore B, Shultz B, Haver DW. Factors contributing to paramedic onscene time during evaluation and management of blunt trauma. *Am J Emerg Med*. 1988;6:443–448.
- Henderson RA, Thomson DP, Bahrs BA, Norman MP. Unnecessary intravenous access in the emergency setting. *Prehosp Emerg Care*. 1998;2:312–316.
- Honigman B, Rohweder K, Moore EE, Lowenstein SR, Pons PT. Prehospital advanced trauma life support for penetrating cardiac wounds. *Ann Emerg Med*. 1990;19:145–150.
- Jones SE, Nesper TP, Alcouloumre E. Prehospital intravenous line placement: a prospective study. *Ann Emerg Med*. 1989;18:244–246.
- Minville V, Pianezza A, Asehnoun K, Cabardis S, Smail N. Prehospital intravenous line placement assessment in the French emergency system: a prospective study. *Eur J Anaesthesiol*. 2006;23:594–597.
- O'Gorman M, Trabulsky P, Pilcher DB. Zero-time prehospital i.v. *J Trauma*. 1989;29:84–86.
- Pace SA, Fuller FP, Dahlgren TJ. Paramedic decisions with placement of out-of-hospital intravenous lines. *Am J Emerg Med*. 1999;17:544–547.
- Pons PT, Moore EE, Cusick JM, Brunko M, Antuna B, Owens L. Prehospital venous access in an urban paramedic system—a prospective on-scene analysis. *J Trauma*. 1988;28:1460–1463.
- Sampalis JS, Tamim H, Denis R, et al. Ineffectiveness of on-site intravenous lines: is prehospital time the culprit? *J Trauma*. 1997;43:608–615.
- Seamon MJ, Fisher CA, Gaughan J, et al. Prehospital procedures before emergency department thoracotomy: “scoop and run” saves lives. *J Trauma*. 2007;63:113–120.
- Slovic CM, Herr EW, Londorf D, Little TD, Alexander BR, Guthmann RJ. Success rates for initiation of intravenous therapy en route by prehospital care providers. *Am J Emerg Med*. 1990;8:305–307.
- Smith JP, Bodai BI, Hill AS, Frey CF. Prehospital stabilization of critically injured patients: a failed concept. *J Trauma*. 1985;25:65–70.
- Spaite DW, Tse DJ, Valenzuela TD, et al. The impact of injury severity and prehospital procedures on scene time in victims of major trauma. *Ann Emerg Med*. 1991;20:1299–1305.
- Benumof JL, Wyte SR, Rogers SN. A large catheter sheath introducer with an increased side-port functional gauge. *Crit Care Med*. 1983;11:660–662.
- Boyle MF, Kuntz B. Saline locks in prehospital care. *Prehosp Disaster Med*. 1994;9:190–192.
- Carducci B, Stein K. Intravenous maintenance with a saline lock intermittent infusion device in the prehospital environment. *Prehosp Disaster Med*. 1994;9:67–70.
- Dalton AM. Prehospital intravenous fluid replacement in trauma: an outmoded concept? *J R Soc Med*. 1995;88:213P–216P.
- Dula DJ, Wood GC, Rejmer AR, Starr M, Leicht M. Use of prehospital fluids in hypotensive blunt trauma patients. *Prehosp Emerg Care*. 2002;6:417–420.
- Eckstein M, Chan L, Schneir A, Palmer R. Effect of prehospital advanced life support on outcomes of major trauma patients. *J Trauma*. 2000;48:643–648.
- Frascone RJ, Jensen JP, Kaye K, Salzman JG. Consecutive field trials using two different intraosseous devices. *Prehosp Emerg Care*. 2007;11:164–171.
- Guisto JA, Iserson KV. The feasibility of 12-gauge intravenous catheter use in the prehospital setting. *J Emerg Med*. 1990;8:173–176.
- Herron H, Falcone R, Dean B, et al. 8.5 French peripheral intravenous access during air medical transport of the injured patient. *Air Med J*. 1997;16:7–10.
- Holcomb JB. Fluid resuscitation in modern combat casualty care: lessons learned from Somalia. *J Trauma*. 2003;54:S46–S51.
- Kaweski SM, Sise MJ, Virgilio RW. The effect of prehospital fluids on survival in trauma patients. *J Trauma*. 1990;30:1215–1218.
- Revell M, Porter K, Greaves I. Fluid resuscitation in prehospital trauma care: a consensus view. *Emerg Med J*. 2002;19:494–498.
- Talving P, Palstedt J, Riddez L. Prehospital management and fluid resuscitation in hypotensive trauma patients admitted to Karolinska University Hospital in Stockholm. *Prehosp Disaster Med*. 2005;20:228–234.
- Turner J, Nicholl J, Webber L, Cox H, Dixon S, Yates D. A randomised controlled trial of prehospital intravenous fluid replacement therapy in serious trauma. *Health Technol Assess*. 2000;4:1–57.

43. Westfall MD, Price KR, Lambert M, et al. Intravenous access in the critically ill trauma patient: a multicentered, prospective, randomized trial of saphenous cutdown and percutaneous femoral access. *Ann Emerg Med.* 1994;23:541–545.
44. Barkana Y, Stein M, Maor R, Lynn M, Eldad A. Prehospital blood transfusion in prolonged evacuation. *J Trauma.* 1999;46:176–180.
45. Cooper DJ, Myles PS, McDermott FT, et al. Prehospital hypertonic saline resuscitation of patients with hypotension and severe traumatic brain injury: a randomized controlled trial. *JAMA.* 2004;291:1350–1357.
46. Holcroft JW, Vassar MJ, Turner JE, et al. 3% NaCl and 7.5% NaCl/dextran 70 in the resuscitation of severely injured patients. *Ann Surg.* 1987;206:279–288.
47. Maningas PA, Mattox KL, Pepe PE, Jones RL, Feliciano DV, Burch JM. Hypertonic saline-dextran solutions for the prehospital management of traumatic hypotension. *Am J Surg.* 1989;157:528–533.
48. Mattox KL, Maningas PA, Moore EE, et al. Prehospital hypertonic saline/dextran infusion for post-traumatic hypotension. The U.S.A. Multicenter Trial. *Ann Surg.* 1991;213:482–491.
49. Sumida MP, Quinn K, Lewis PL, et al. Prehospital blood transfusion versus crystalloid alone in the air medical transport of trauma patients. *Air Med J.* 2000;19:140–143.
50. Vassar MJ, Fischer RP, O'Brien PE, et al. A multicenter trial for resuscitation of injured patients with 7.5% sodium chloride. The effect of added dextran 70. The Multicenter Group for the Study of Hypertonic Saline in Trauma Patients. *Arch Surg.* 1993;128:1003–1011.
51. Vassar MJ, Perry CA, Gannaway WL, Holcroft JW. 7.5% sodium chloride/dextran for resuscitation of trauma patients undergoing helicopter transport. *Arch Surg.* 1991;126:1065–1072.
52. Vassar MJ, Perry CA, Holcroft JW. Prehospital resuscitation of hypotensive trauma patients with 7.5% NaCl versus 7.5% NaCl with added dextran: a controlled trial. *J Trauma.* 1993;34:622–632.
53. Branas CC, Sing RF, Davidson SJ. Urban trauma transport of assaulted patients using nonmedical personnel. *Acad Emerg Med.* 1995;2:486–493.
54. Rosa DM, Griffin CC, Flanagan JJ Jr, Machiedo GW. A comparison of intravenous access sites for bolus injections during shock and resuscitation after emergency room thoracotomy with and without aortic crossclamping. *Am Surg.* 1990;56:566–570.
55. Fowler R, Gallagher JV, Isaacs SM, Ossman E, Pepe P, Wayne M. The role of intraosseous vascular access in the out-of-hospital environment (resource document to NAEMSP position statement). *Prehosp Emerg Care.* 2007;11:63–66.
56. Von Hoff DD, Kuhn JG, Burriss HA III, Miller LJ. Does intraosseous equal intravenous? A pharmacokinetic study. *Am J Emerg Med.* 2008;26:31–38.
57. Hashiguchi N, Lum L, Romeril E, et al. Hypertonic saline resuscitation: efficacy may require early treatment in severely injured patients. *J Trauma.* 2007;62:299–306.
58. Holcroft JW, Vassar MJ, Perry CA, Gannaway WL, Kramer GC. Use of a 7.5% NaCl/6% Dextran 70 solution in the resuscitation of injured patients in the emergency room. *Prog Clin Biol Res.* 1989;299:331–338.
59. Maningas PA. Resuscitation with 7.5% NaCl in 6% dextran-70 during hemorrhagic shock in swine: effects on organ blood flow. *Crit Care Med.* 1987;15:1121–1126.
60. Vassar MJ, Perry CA, Holcroft JW. Analysis of potential risks associated with 7.5% sodium chloride resuscitation of traumatic shock. *Arch Surg.* 1990;125:1309–1315.
61. Wade CE, Dubick MA, Vassar MJ, Perry CA, Holcroft JW. Plasma dextran concentrations in trauma patients administered hypertonic saline-dextran-70. *Clin Chem.* 1996;42:779–780.
62. Pinto FC, Capone-Neto A, Prist R, E Silva MR, Poli-de-Figueiredo LF. Volume replacement with lactated Ringer's or 3% hypertonic saline solution during combined experimental hemorrhagic shock and traumatic brain injury. *J Trauma.* 2006;60:758–763.
63. Stern SA, Kowalenko T, Younger J, Wang X, Dronen SC. Comparison of the effects of bolus vs. slow infusion of 7.5% NaCl/6% dextran-70 in a model of near-lethal uncontrolled hemorrhage. *Shock.* 2000;14:616–622.
64. Watters JM, Tieu BH, Differding JA, Muller PJ, Schreiber MA. A single bolus of 3% hypertonic saline with 6% dextran provides optimal initial resuscitation after uncontrolled hemorrhagic shock. *J Trauma.* 2006;61:75–81.
65. Schmand JF, Ayala A, Morrison MH, Chaudry IH. Effects of hydroxyethyl starch after trauma-hemorrhagic shock: restoration of macrophage integrity and prevention of increased circulating interleukin-6 levels. *Crit Care Med.* 1995;23:806–814.
66. Schmand JF, Ayala A, Chaudry IH. Effects of trauma, duration of hypotension, and resuscitation regimen on cellular immunity after hemorrhagic shock. *Crit Care Med.* 1994;22:1076–1083.
67. Gould SA, Moore EE, Hoyt DB, et al. The first randomized trial of human polymerized hemoglobin as a blood substitute in acute trauma and emergent surgery. *J Am Coll Surg.* 1998;187:113–120.
68. Gould SA, Moore EE, Hoyt DB, et al. The life-sustaining capacity of human polymerized hemoglobin when red cells might be unavailable. *J Am Coll Surg.* 2002;195:445–452.
69. Krausz MM, Bashenko Y, Hirsh M. Crystalloid and colloid resuscitation of uncontrolled hemorrhagic shock following massive splenic injury. *Shock.* 2001;16:383–388.
70. Sumas ME, Legos JJ, Nathan D, Lamperti AA, Tuma RF, Young WF. Tonicity of resuscitative fluids influences outcome after spinal cord injury. *Neurosurgery.* 2001;48:167–172.
71. Varicoda EY, Poli de Figueiredo LF, Cruz RJ Jr, Silva LE, Rocha e Silva M. Blood loss after fluid resuscitation with isotonic or hypertonic saline for the initial treatment of uncontrolled hemorrhage induced by spleen rupture. *J Trauma.* 2003;55:112–117.
72. Krausz MM, Hirsh M. Bolus versus continuous fluid resuscitation and splenectomy for treatment of uncontrolled hemorrhagic shock after massive splenic injury. *J Trauma.* 2003;55:62–68.
73. Hambly PR, Dutton RP. Excess mortality associated with the use of a rapid infusion system at a level 1 trauma center. *Resuscitation.* 1996;31:127–133.