

# **PRACTICE MANAGEMENT GUIDELINES FOR GERIATRIC TRAUMA**

The EAST Practice Management Guidelines Work Group:

David G. Jacobs, MD, **Committee Co-Chair**  
Carolinas Medical Center  
Charlotte, NC

Brian Ray Plaisier, MD, **Committee Co-Chair**  
Bronson Hospital  
Kalamazoo, MI

Philip S. Barie, MD  
NYH – Cornell Medical Center  
New York, NY

Jeffrey S. Hammond, MD  
Robert Wood Johnson Medical School  
New Brunswick, NJ

Michele R. Holevar, MD  
Mt. Sinai Hospital  
Chicago, IL

Karlene E. Sinclair, MD  
Morehouse School of Medicine  
Atlanta, GA

Thomas M. Scalea, MD  
University of Maryland Medical Center  
Baltimore, MD

Wendy Wahl, MD  
University of Michigan Health System  
Ann Arbor, MI

## INTRODUCTION

Advanced age is a well-recognized risk factor for adverse outcomes following trauma. A substantial body of literature, much of it cited within this document, demonstrates increased morbidity and mortality in geriatric trauma patients compared to their younger counterparts. Whether this outcome difference is due to the decreased physiologic reserve that accompanies aging, a higher incidence of pre-existing medical conditions in the geriatric patient, or to other factors yet to be identified, remains unclear. It is clear, however, that good outcomes can be achieved in this patient population when appropriately aggressive trauma care is directed towards geriatric patients with survivable injuries. Implicit in the above statement is the need to identify, as soon as possible following injury, those patients who will benefit from aggressive resuscitation, timely injury management, and post-trauma rehabilitation. It is equally important, however, to limit these intensive and expensive treatment modalities to patients whose injuries are not only survivable, but are compatible with an acceptable quality of life.

Our purpose in developing this guideline was to provide the trauma practitioner with some *evidence-based* recommendations that could be used to guide decision-making in the care of the geriatric trauma patient. We began this process by first developing a series of questions, the answers to which we hoped could be supported by the existing scientific literature. The initial set of questions were as follows:

1. Is age itself a marker of increased morbidity/mortality? If so, what age should be used?
2. Is age instead a surrogate for increased pre-existing conditions (PEC's)? If so, which pre-morbid conditions are particularly predictive of poor outcomes?
3. Should age itself be a criterion for triage from the field directly to a trauma center, regardless of Glasgow Coma Scale (GCS) score, trauma score (TS), etc.? If so, what age should be used?
4. Do trauma centers have better outcomes with geriatric trauma than non-trauma centers?
5. Are there specific injuries, scores [Injury Severity Score (ISS), TS, GCS, etc], or PEC/age combinations in geriatric trauma patients that are so unlikely to be survivable that a non-aggressive approach from the outset could be justified?
6. What resuscitation end-points should be used for the geriatric trauma patient?
7. Should *all* geriatric trauma patients receive invasive hemodynamic monitoring? If so, what specific types of monitoring should be used? If not, which geriatric patients benefit from invasive monitoring?
8. Are there specific types of therapies that should be employed routinely in trauma patients (e.g.  $\beta$ - blockade, nitroglycerin infusions, etc.)?
9. How are outcomes measured in geriatric trauma? Which specific outcome measures should be used?

Unfortunately, after examining the available literature, it is clear that *evidence-based* responses to all of the questions raised above are not possible. As the accompanying evidentiary tables demonstrate, there are a few, if any, prospective, randomized, controlled trials which definitively address any of the above issues. Secondly, there is a lack of uniformity as to a specific age criterion for geriatric trauma. As shown in the accompanying evidentiary table, geriatric trauma is variously defined in the literature as age greater than or equal to 55, 60, 65, 70, 75, and even 80 years of age. There is even literature support for increased mortality from trauma beginning at age 45! Furthermore, since age is a continuous variable, and not a dichotomous one, adverse outcomes associated with geriatric trauma are likely to increase in a

continuous fashion which age as opposed to a stepwise leap as a given patient reaches a specific age. Third, there is no concise definition of a geriatric trauma patient. In some studies, all patients over a given age are included, whereas in others, patients with penetrating injuries, burns, and those with minor injuries, such as slip-and-falls, are excluded. Some studies include all patients regardless of hemodynamic instability or injury severity, while others impose strict entrance criteria or exclude patients who do not survive for a predetermined period of time following admission. Such lack of uniformity with regards to inclusion criteria makes it difficult to compare outcomes across different patient populations. Finally, much of the literature concerning geriatric trauma is relatively “old”, that is, published more than 10 years ago. Given the significant improvements in patient care which have occurred over the past 10 to 20 years, recommendations based upon outcomes achieved more than 10 years ago may not be applicable to today’s geriatric trauma patient.

Despite the above-mentioned shortcomings, our committee still felt that it was important to summarize the available literature and make evidence-based recommendations where satisfactory evidence did exist. In light of the nine questions raised above, three broad areas of focus emerged within this guideline: Issues of Geriatric Trauma Triage, Issues of Geriatric Trauma Resuscitation, and Issues of Outcome Measurement in Geriatric Trauma. Although there was considerable overlap among these three areas, each issue has been addressed separately within this guideline and, accordingly, three separate “sub-guidelines,” each with its own recommendations, evidentiary table, and areas for future research, comprise this practice management guideline for geriatric trauma. It is hoped that the information provided within these three sub-guidelines, will provide evidence-based support for the difficult decisions which are required to achieve optimal outcomes in this difficult, but ever increasing, patient group.

## **TRIAGE ISSUES IN GERIATRIC TRAUMA**

### **I. STATEMENT OF THE PROBLEM**

The process of triage, as it relates to the geriatric trauma patient, is an attempt to provide the patient with the appropriate intensity of medical resources, taking into account the severity of illness, the cost and availability of medical resources, the prognosis for functional survival and, if known, the expressed desires of the patient. For the geriatric trauma patient this process begins in the pre-hospital phase of care where decisions must be made regarding the appropriate patient destination, trauma center versus non-trauma center. In the resuscitative phase of trauma care, triage decisions regarding patient destination must again be made, specifically whether patient circumstances dictate provision of intensive care resources or whether standard trauma inpatient care will suffice. Throughout the hospital phase of care, the patients must be “triaged” towards or away from operative procedures, invasive and expensive critical care therapies, and powerful, yet potentially dangerous pharmacologic treatment options, decisions which, again, must be based upon the likelihood of achieving a good, long-term outcome for the patient. An increasingly common circumstance, particularly in the geriatric trauma patient, involves the decision to withdraw, or perhaps not even institute, an aggressive course of treatment, when the clinical circumstances are incompatible with a quality of life which all parties concerned would deem acceptable. Fundamental to all of these “triage” decisions, is the ability to predict with reasonable accuracy, what a particular patient’s outcome might be depending on which “triage” decision is made. In order to be of any value to the trauma practitioner, and ultimately to the patient and his or her family, the clinical variables upon which these predictions are to be made must be easy to obtain, reliable, and available to the trauma practitioner within a relatively short period of time following injury. The task, therefore, of this particular subcommittee was to determine whether there existed adequate support in the scientific literature to develop recommendations regarding 1) appropriate criteria for triage of the geriatric trauma patient to trauma centers, 2) the clinical variables which would be useful in predicting the need for intensive care resources for the geriatric trauma patient, and 3) those clinical circumstances where a non-aggressive approach from the outset could be justified.

### **II. PROCESS**

An initial computerized search was undertaken using Medline with citations published between the years of 1966 and 1999. Using the search words “geriatric”, “trauma”, “elderly”, and “injury”, and by limiting the search to citations dealing with human subjects and published in the English language, well over 2,300 citations were identified. From this number were then excluded letters to the editor, case reports, reviews, and a large number of articles dealing with minor injury mechanisms, particularly hip fractures from slip-and-falls. An additional cause for exclusion of references was publication prior to 1975 as it was felt that the trauma care provided at this time was so different compared to current trauma care that recommendations based upon data from this earlier time period would not be valid. The abstracts of the remaining citations were each reviewed, and those articles that did not address prognostic variables or other issues pertinent to the triage of the geriatric trauma patient were further excluded. This yielded a total of 32 articles that comprised the initial evidentiary table. The bibliographies of these 32 articles were then further reviewed and additional 13 articles meeting the above-mentioned criteria were

added for a total of 45 references within the evidentiary table. Each reference was then reviewed by three trauma surgeons, and consensus reached regarding appropriate classification of each reference according to the Canadian and United States Preventive Task Force.

Criteria for achieving a specific classification and the number of articles for each class are shown below:

**Class I:** Prospective randomized controlled trials - the gold standard of clinical trials. Some may be poorly designed, have inadequate numbers, or suffer from other methodological inadequacies. (**0 references**)

**Class II:** Clinical Studies in which data was collected prospectively, and retrospective analyses that were based on clearly reliable data. Types of studies so classified include observational studies, cohort studies, prevalence studies and case control studies. (**2 references**)

**Class III:** Studies based on retrospectively collected data. Evidence used in this class indicate clinical series, database or registry review, large series of case reviews, and expert opinion (**44 references**)

(The total number of classified references exceeds the total number of references by one because one two-part study was classified as both a Class II and Class III reference.)

### III. RECOMMENDATIONS

#### A. Level I

There is insufficient Class I and Class II data to support any standards regarding triage of geriatric trauma patients.

#### B. Level II

1. Advanced patient age should lower the threshold for field triage directly to a trauma center.

#### C. Level III

1. All other factors being equal, advanced patient age, in and of itself, is not predictive of poor outcomes following trauma, and therefore should NOT be used as the sole criterion for denying or limiting care in this patient population.
2. The presence of pre-existing medical conditions (PEC's) in elderly trauma patients adversely affects outcome. However this effect becomes progressively less pronounced with advancing age.
3. In patients 65 years of age and older, a  $GCS \leq 8$  is associated with a dismal prognosis. If substantial improvement in GCS is not realized within 72 hours of injury, consideration should be given to limiting further aggressive therapeutic interventions.
4. Post-injury complications in the elderly trauma patient negatively impact survival and contribute to longer lengths of stay in survivors and non-survivors compared to younger trauma patients. Specific therapies designed to prevent and/or reduce the occurrence of complications (particularly iatrogenic complications) should lead to optimal outcomes in this patient population.

5. With the exception of patients who are moribund on arrival, an initial aggressive approach should be pursued with the elderly trauma patient, as the majority will return home, and up to 85% will return to independent function.
6. In patients 55 years of age and older, an admission base deficit  $\leq 6$  is associated with a 66% mortality. Patients in this category may benefit from in-patient triage to a high-acuity nursing unit.
7. In patients 65 years of age and older, a Trauma Score  $< 7$  is associated with a 100 % mortality. Consideration should be given to limiting aggressive therapeutic interventions.
8. In patients 65 years of age and older, an admission respiratory rate  $< 10$  is associated with a 100 % mortality. Consideration should be given to limiting aggressive therapeutic interventions.
9. Compared to younger trauma patients, patients 55 years of age and older are at considerably increased risk for undertriage to trauma centers even when these older patients satisfy appropriate triage criteria. The factors responsible for this phenomenon must be identified and strategies developed to counteract it.

#### IV. SCIENTIFIC FOUNDATION

Triage is the process whereby the patient's medical needs are matched with the available medical resources. For the geriatric trauma patient, the process begins in the pre-hospital arena, where pre-hospital providers must decide on the basis of relatively scant clinical information whether a patient should bypass the local hospital in favor of a trauma center. The American College of Surgeons Committee on Trauma (ACS-COT), among other medical organizations, in its manual, "Optimal Resources for the Care of the Trauma Patient" has published a set of triage criteria to aid pre-hospital providers in identifying appropriate patients for direct transport to trauma centers<sup>1</sup>. Within this document, it is suggested that patients greater than age 55 should be "considered" for direct transport to a trauma center, apparently without regard to the severity of injury. This recommendation is based upon a substantial medical literature that demonstrates significantly worse outcomes for geriatric trauma patients compared to their non-geriatric counterparts. One of the earliest studies to look at the influence of age on outcome from major trauma was the Major Trauma Outcome Study, sponsored by the ACS-COT. Data from 3,833 patients 65 years and older was compared to that of 42,944 patients less than 65 years of age. Mortality rose sharply between age 45 and 55 and doubled at age 75 years. This age-dependent survival decrement occurred at all ISS values, for all mechanisms of injury, and for all body region.<sup>2</sup> Numerous other studies have supported the findings that the effect of trauma on the elderly is more serious than that on younger patients.<sup>3-9</sup>

Given these findings, some authors have suggested triaging elderly trauma victims to trauma centers at a much lower threshold than similarly injured younger patients, in order to minimize mortality and morbidity.<sup>10</sup> Support for this recommendation can be found in a study by Smith et al, documenting fewer complications for elderly femur fracture patients treated at trauma centers Vs non-trauma centers.<sup>11</sup> In spite of these poorer outcomes, trauma patients 55 years of age and older are frequently triaged to non-trauma hospitals even when they satisfy well-defined anatomic or physiologic criteria. Compliance with physiologic criteria appears to be especially troublesome.<sup>12</sup> In two unrelated studies, undertriage in patients over the age of 55 was *twice*

that of younger patients,<sup>12, 13</sup> while a similar study demonstrated even worse results for patients over the age of 65.<sup>14</sup>

The factors responsible for the increased morbidity and mortality seen in geriatric trauma are not entirely clear. It has been suggested that it is not patient age per se, but the high incidence of pre-existing medical conditions in the geriatric patient that accounts for the difference. Others have suggested that the elderly, simply by virtue of being more frail, sustain a greater degree of injury in response to a given impact, compared to their younger counterparts. The existing medical literature was therefore reviewed in an attempt to identify clinical factors that might be used to triage geriatric trauma patients to either aggressive Vs non-aggressive treatment strategies.

## **AGE AND OUTCOME**

It is difficult to find consensus in the existing literature regarding the relationship between the patient age and outcome. Many of the reasons for this failure have been mentioned above, and include differences in the age definition of geriatric trauma, and differences in inclusion criteria for the various studies. In addition to these two factors, there is a lack of uniformity regarding the length of follow-up required to define a poor outcome. This has been variably defined as death within 24 or 48 hours of injury, death prior to ICU or hospital discharge, and even death/vegetative outcome at three or four years post injury. Furthermore, there are wide variances in the statistical methods used to explore the relationship between age and outcome. Many authors have documented a statistically significant difference between the mean age of geriatric survivors compared with the mean age of geriatric *non*-survivors, and thus have concluded that age is significantly *associated* with poor outcome. Other authors have applied logistic regression analysis to their data set to determine which particular factors are *predictive* of adverse outcomes. Given the wide variation in inclusion criteria, outcome variables and statistical methods present within the existing literature, the conflicting results regarding age and outcome are not surprising.

Certainly, the largest data set examined to date is that published by Morris et al in 1990. These authors examined 199,737 trauma admissions, aged 15 and older, to acute care hospitals in the state of California during 1986. Using logistic regression techniques, the Injury Severity Score (ISS) was found to be the best predictor of mortality in trauma patients, but age, gender, and pre-existing medical conditions (PEC'S) were also found to be *independent* predictive factors of mortality. Mortality was defined as in-hospital death. The authors also found that while the mortality from minor injury (ISS <9) begins to increase beyond the age of 65, the mortality for moderate injuries (ISS 9-24) begins to increase at 45 years of age.<sup>4</sup> This increase in trauma mortality beginning at age 45 had been confirmed by other investigators as well.<sup>2, 10, 15</sup> Several authors have examined the relationship between *in-hospital* mortality and age, with differing conclusions. Pelicane et al demonstrated a statistically significant difference in age between elderly non-survivors and elderly survivors in a series of 374 geriatric trauma patients, defined as age greater than or equal to 65 years. Five of the deaths in this series occurred in the Emergency Department, a subset of patients that has been excluded from the analyses in other series. Burn patients, however, were excluded from this series.<sup>16</sup> In a similar study, performed by Osler, of 100 geriatric trauma patients 65 years or older, no significant differences in age was found between elderly survivors and non-survivors.<sup>9</sup> Despite the fact that patients who expired prior to transfer to the operating room or to the ICU were excluded in Osler's series, mortality in

this series was more than twice that in Pelicane's series. Perhaps this is explained by the lower mean Trauma Score (TS) in Pelicane's series relative to that of Osler (13 versus 15.4). Pelicane's series contains nearly four times as many patients which raises, the possibility of a type 2 statistical error with regards to Osler's inability to demonstrate a statistical difference between the ages of geriatric trauma survivors and non-survivors. A large and more recent study of 448 patients, 65 years and older, employed a logistic regression analysis and demonstrated age to be significantly predictive of both early (<24 hours) and late (>24 hours) mortality.<sup>17</sup> In this analysis, survival was used as the outcome variable, with "geriatric status" (age greater than or equal to 65) entered into the logistic regression equation. In so doing, "geriatric status" was associated with a 2.46-fold increased likelihood of early mortality and a 4.64-fold increased risk of late mortality. However, an even larger study yet, consisting of 852 patients, reported on by Knudson et al, using stepwise discriminant analysis, did not find age to be predictive of in-hospital death. The authors reported a 1.33-fold increased risk of death associated with age status greater than 75 years, just barely missing statistical significance with a p-value of 0.06. Interestingly enough, however, the age of 75 years was entered into the discriminant analysis, not the age of 65 or greater which was the authors' original age definition for entrance into the study. Perhaps statistical significance would have been demonstrated had age 65 or greater been used in the discriminant analysis.<sup>18</sup>

Two studies specifically examined the relationship between age and in-hospital mortality for geriatric trauma patients admitted to the Intensive Care Unit. Neither found any association between age and outcome. In a small series of 39 patients requiring intensive care unit admission and placement of pulmonary and radial artery catheters, Horst et al reported no significant difference in age between elderly survivors and non-survivals. As would be expected, overall mortality (31%) was high in this intensive care population of patients greater than 60 years of age. Logistic regression analysis was not performed in this study, probably due to the overall low number of patients.<sup>19</sup> A more recent study by Shabot and colleagues examined two subsets of geriatric trauma patients, those between the ages of 65 and 74, and those 75 years and older. Outcomes in these 99 geriatric trauma patients were then compared to 940 "younger" patients between the ages of 13 and 64, all of who were admitted to a surgical intensive care unit (SICU). SICU mortality was then examined by comparing survivors with non-survivors, regardless of age. As would be expected, there was no significant difference in age between non-survivors and survivors (39.0 years versus 34.8 years), likely due to the tenfold larger number of patients seen in the "younger" patient group.<sup>20</sup>

Finally, several studies have examined the relationship of age to more long-term outcomes, although no clear consensus is evident. DeMaria et al studied a group of 82 trauma patients over the age of 65 years. Patients with penetrating injury and isolated orthopedic injury were excluded, as were patients sustaining thermal injury. Survival was defined at six months post-injury. Not only were non-survivors older, but they also demonstrated higher ISS's and more complications. Based on these findings, the authors developed the Geriatric Trauma Survival Score (GTSS), and then prospectively tested it on 61 patients, with 92% accuracy. Unfortunately, the GTSS, though perhaps accurate, has little triage value at the time of patient admission, as it requires information not available to the practitioner at that time.<sup>6</sup> Van Der Sluis compared early and late mortality between elderly trauma patients and elderly hip fracture patients. Early mortality was higher for the trauma patients, but survival seven to eight years following injury was similar between the two groups. Logistic regression analysis was employed to identify predictors of late mortality, and demonstrated age to be a significant predictor.<sup>21</sup> van

Aalst, in a study of blunt geriatric injury with a mean follow-up of almost 3 years, employed logistic regression analysis to demonstrate an association between poor outcome and age > 75.<sup>5</sup> Oreskovich, however, failed to demonstrate any relationship between age and outcome at one year following injury in a group of 100 patients age 70 and greater.<sup>22</sup> Broos, in two separate publications examining six-month outcome in trauma patients aged 65 and greater, did not find age to be predictive of mortality.<sup>23, 24</sup> Inclusion criteria for each of these 3 latter studies were vaguely defined, and Broos' 18% mortality is inexplicably low compared to other series of similarly injured patients.<sup>9, 17, 25, 26</sup> A larger study, with a more plausible mortality, was published by Battistella et al in 1998. This study involved 279 geriatric trauma patients, which the authors defined as age greater than 75. Mean ISS in this patient group was 9.4, and associated mortality was 23%. Using logistic regression analysis, the authors found that poor outcome, defined in this study as survival less than six months following *hospital discharge*, was not predicted by patient age.<sup>27</sup> The issue of long-term survival and quality of life in the geriatric trauma patient is discussed more fully below.

Can the age of a geriatric patient, then, be used to predict outcome following trauma? While age appears to have some value in mortality projections for a *population* of geriatric trauma patients, there is certainly no literature support for a specific age above which geriatric trauma *in-hospital mortality* can be predicted with any degree of confidence. It has been suggested, however, that *early* mortality may not be the best outcome measure in geriatric trauma, due to a high percentage of poor long-term *functional* survival in elderly trauma patients surviving hospital discharge.<sup>22</sup> The preponderance of available literature, however, suggest more favorable long-term outcomes, with up to 85% of survivors functioning independently at home at follow-up intervals as long as six years post injury.<sup>5, 27-31</sup> Thus, given reasonable long-term functional outcomes for geriatric trauma patients surviving hospitalization, and the inability of patient age, by itself, to predict in-hospital mortality, advanced patient age should not be used as the sole criteria for denying or limiting care in the geriatric trauma population.

### **PRE-EXISTING CONDITIONS (PEC'S) AND OUTCOME**

If *chronologic* age, then, is not useful in predicting geriatric trauma survival, perhaps it is the patient's *physiologic* age, or the nature and extent of pre-existing medical conditions (PEC's), that determines outcome. Since the frequency of PEC's does increase with age, it may be difficult to separate these two factors and their relationships to adverse outcomes in geriatric trauma. Unfortunately, due, once again, to a wide variety of age definitions for geriatric trauma, statistical methodologies, and outcome measures, the literature addressing the prognostic value of PEC's in geriatric trauma outcome is inconclusive. The largest studies, and those with the best statistical methodology, do seem to demonstrate a significant predictive capacity of PEC's for adverse outcomes in geriatric trauma. Morris et al, in two separate publications in 1990, examined hospital discharge data for trauma patients in California for the year 1986. Using logistic regression analysis in both studies, Morris was able to demonstrate that PEC's were important predictive factors of mortality, independent of age. The effect of PEC's on mortality, however, became less important in patients over the age of 65, perhaps because at this age, chronologic age becomes the predominant predictor of mortality, and the added presence of PEC's does little to increase trauma mortality further.<sup>4, 32</sup> Similarly, Milzman et al, in the study of nearly 8,000 trauma patients, noted a threefold increase in trauma mortality in patients with PEC's, compared to those without. Once again, the effect of PEC's on mortality was noted to be independent of age although, like Morris et al, these authors noted a decreasing influence of

PEC's on trauma mortality with advanced age.<sup>33</sup> A more recent study published in 1997 by Gubler et al, examined risk factors for mortality among a group of 9,424 trauma patients, aged 67 and greater, who were discharged from acute care hospitals within the state of Washington in 1987. For each trauma patient in the series, four uninjured patients, matched for age and gender, were identified from the same HCFA database. Co-morbid diagnoses (PEC's) were identified for each patient, and a Co-Morbid Diagnosis Index Score calculated. This score is a weighted index which takes into account not only the number, but also the severity of PEC's.<sup>34</sup> Using Cox proportional hazards regression, Gubler found that patients with PEC's were anywhere between 2.0 and 8.4 times as likely to die within five years of injury compared to those without PEC's, depending upon the number and severity of PEC's.<sup>8</sup> Several smaller studies, each reporting the experience of a single trauma center, and using logistic regression analysis, confirmed the value of PEC's as predictive factors of poor outcome in geriatric trauma, although inclusion criteria, and age and outcome definitions were not uniform among these studies.<sup>17, 27, 35</sup> Other studies have refuted these findings, but suffer from some methodological and statistical shortcomings that weaken their conclusions.<sup>19, 22-24, 36, 37</sup>

### SEVERITY OF INJURY SCORING AND OUTCOME

A number of physiologic and anatomic "scores" have been shown to correlate with geriatric outcome. These include Trauma Score (TS), Revised Trauma Score (RTS), Glasgow Coma Scale (GCS), Acute Physiology and Chronic Health Evaluation Score (APACHE), Acute Physiologic Score (APS), Simplified Acute Physiology Score (SAPS), Injury Severity Score (ISS), Maximal Abbreviated Injury Score (MAIS), and the Geriatric Trauma Survival Score (GTSS). In addition, although not "scores" in the typical sense, geriatric trauma outcome has also been correlated with initial blood pressure, respiratory rate, and base deficit. Although most, if not all, of these scores do correlate with geriatric outcome, from the perspective of field or emergency department triage, many of these scores have little value in that they are not derivable at the moment that these particular triage decisions need to be made. This would apply particularly to APACHE, APS, SAPS, MAIS, ISS and GTSS. These scores, however, perhaps in combination with patient age, may have some value in the prediction of lethal outcomes in geriatric trauma, and, therefore, may be valuable triage tools in the Intensive Care Unit. These scores will therefore be discussed solely within that context below.

On the other hand, measures of *physiologic* derangement, whether obtained via physical examination or chemical analysis, may help to identify patients who will perhaps benefit from aggressive resuscitation strategies (and should therefore be triaged to an intensive care unit), as well as those where further resuscitated efforts are futile (thus facilitating earlier termination of resuscitation). TS (or RTS) and its components (blood pressure, respiratory rate and GCS) are the most readily obtainable, objective physiologic data available either to the pre-hospital provider or to the trauma resuscitation team in the Emergency Department. The prognostic value of each of these variables as they relate to geriatric trauma outcome will be discussed further below. (The prognostic value of GCS will be discussed within the section entitled "Outcome from Geriatric Head Injury.") Of those chemical analyses available in the Emergency Department, only base deficit has been subjected to sufficient scientific study, and is sufficiently relevant to geriatric trauma resuscitation that it can be included within the discussion below of potentially useful triage "scores."

The TS assesses five physiologic functions (blood pressure, respiratory rate, respiratory effort, GCS and capillary refill), yielding a minimal score of zero and a maximal score of 16. The RTS is a simplified version of the TS, which deletes the assessment of respiratory effort and capillary refill, resulting in a range of scores between zero and eight. Several studies have demonstrated the predictive value of TS (and RTS) on geriatric trauma mortality, although a specific numerical score signifying a fatal injury has not identified by these authors.<sup>17, 20</sup> Horst was unable to find any relationship between TS and geriatric trauma patient mortality, although the small number of patients and the rather narrow entrance criteria of the study limit the applicability of this finding.<sup>19</sup> Two other studies, however, suggest that TS may be a useful triage tool in the early stages following geriatric trauma. In a case-matched review of 100 patients age 65 and above that suffered injuries severe enough to necessitate hospitalization, no elderly patient was able to survive a TS < 9. More dramatically, no geriatric patient with a TS < 7 survived long enough to reach the hospital and be included in the study. The authors felt this to be of importance in allowing more realistic counseling of patients and their families.<sup>9</sup> These findings were reinforced by a review from three trauma centers of 852 patients, age 65 and older, in which a TS < 7 was associated with a 100% mortality.<sup>18</sup> These data suggest that aggressive care under these circumstances is likely to be futile, and that consideration should be given to limiting intensive therapy when a geriatric patient presents with a TS < 7. In addition to the prognostic value of TS, Knudson's data also revealed a 100% mortality in patients 65 years and older who presented with a respiratory rate < 10. Here, too, consideration should be given to limiting aggressive therapeutic interventions. In addition to its role in the prediction of fatal outcomes in geriatric trauma patients, TS may also have implications for intensive care unit triage. In a study of 374 patients aged 65 and older, mortality was noted to be only 5% with a TS of 15 or 16, but 25% in patients with a TS of 12 to 14, and 65% in patients with a TS < 12.<sup>16</sup> Thus, patients with TS between 7 and 14 may benefit from aggressive resuscitation strategies and triage to a critical care unit.

Measurement of arterial base deficit may provide useful information regarding the extent of shock and the adequacy of resuscitation in trauma patients, and may therefore be useful in early decision-making and resource allocation. In a series of 274 elderly trauma patients, defined for the purposes of this study as age greater than or equal to 55, arterial blood gases obtained within one hour of a patient admission were correlated with ICU length of stay and mortality. Base deficits were characterized as mild (-3 to -5), moderate (-6 to -9), and severe (? -10). As expected, elderly patients with severe base deficits had a high mortality, 80% in this series. However, geriatric trauma mortality was still markedly elevated at 60% in patients with only moderate base deficits. Even a "normal" base deficit carried a mortality of 24%.<sup>7</sup> Thus, early determination of admission base deficit in geriatric trauma patients may facilitate early identification of "occult shock" and identify a subgroup of patients who may benefit from more intensive monitoring and resuscitation.

Injury Severity Score (ISS) is probably the most widely studied anatomic or physiologic severity of illness score yet to be correlated with geriatric trauma outcome. Most authors have found it to be a strong predictor of outcome in geriatric trauma,<sup>9, 16, 17, 20, 21, 25</sup> while two large studies claimed that it is the best predictor of mortality in geriatric trauma.<sup>4, 18</sup> Others, however, have failed to demonstrate any such relationship.<sup>19, 22-24, 27</sup> Whether or not such a relationship does indeed exist, ISS is severely limited in its prognostic capability due to significant delays in obtaining sufficient data to calculate the score. It has, therefore, probably very little prognostic value in geriatric trauma, and, even then, only in patients in whom the question of futility has

been raised. Despite the abundance of literature examining the relationship between ISS and geriatric trauma patient outcome, only two publications contain any ISS data that might be considered useful prognostically. Van der Sluis reported on a series of 121 trauma patients age 60 and greater, all with ISS  $\leq$  16. No patient with an ISS greater than or equal to 50 survived in this series. The authors, however, do emphasize the importance of not using the ISS to predict outcomes in individual patients.<sup>35</sup> A study by Carrillo, published in 1994, reported on 94 blunt trauma victims age 65 or greater. Mortality correlated well with APACHE II, but the combination of APACHE II and ISS performed better than APACHE II alone. All patients with APACHE II  $\geq$  15 and ISS  $\geq$  30 died, but this accounted for only one-third of all deaths in this series.<sup>28</sup> Thus, it would appear that there is little, if any, support in the literature to justify withdrawal of care based upon any combination of age and ISS. Likewise, for SAPS, APS, and MAIS, there is no literature support for the use of any of these scores to predict individual patient outcome following geriatric trauma.<sup>3, 19, 20, 26</sup> Finally, mention should be made of the GTSS, the Geriatric Trauma Survival Score. This score was derived by DeMaria, based upon his experience with 82 blunt trauma patients over the age of 65 years. The formula to calculate GTSS uses patient age, ISS, and the presence or absence of cardiac and septic complications to predict patient outcome.<sup>6</sup> Given the inadvisability mentioned above of using ISS to predict individual patient outcome, as well as the fact that information regarding the presence or absence of complications will not be obtainable prior to hospital discharge, the GTSS clearly has no role in guiding decision-making, a point which the authors themselves emphasize. Interestingly, a larger and more recent study of blunt trauma patients aged 60 and over, failed to demonstrate any relationship between the GTSS and survival.<sup>25</sup>

## COMPLICATIONS AND OUTCOME

It is generally acknowledged that when the geriatric trauma patient sustains complications during their initial hospitalization that overall outcome is adversely affected. Both DeMaria and Osler, in comparing elderly survivors with non-survivors, have noted a statistically higher incidence of cardiac and septic complications<sup>6</sup> and respiratory complications<sup>9</sup> in non-survivors. Other authors, employing logistic regression statistical methodology, have identified cardiac, infectious, and pulmonary complications as independent predictors of poor outcome following geriatric trauma.<sup>5, 17, 25</sup> In addition to the specific types of complications sustained by the geriatric trauma patient, the *number* of complications sustained by a given geriatric trauma patient has been identified as a risk factor for poor outcomes. Smith, in a study of 456 trauma patients aged 65 and over, reported a 5.4% mortality for those patients with no complications, 8.6% for those with one complication, and 30% for those with more than one complication.<sup>37</sup> Similar results have been noted for geriatric patients sustaining traumatic brain injuries.<sup>15</sup>

In spite of the well documented relationship between complications and outcome in geriatric trauma, triage decisions are rarely, if ever, affected by this information. Early triage decisions, whether in the field or in the emergency department, clearly cannot be based upon the presence or absence of complications which yet to occur. Furthermore, there is no data to suggest that any particular number, or type, of complications will allow identification of the *individual* geriatric trauma patient destined for an outcome so dismal that a non-aggressive course of treatment could be justified. In light of these findings, efforts should be focused on the development and implementation of strategies aimed at the *prevention* of complications in the geriatric trauma patient. The importance of complication prevention is highlighted in a study by Pellicane which

revealed that preventable complications contributed to mortality in 32% of all deaths in this series and in 62% of deaths related to multiple organ systems failure.<sup>16</sup>

## OUTCOME FROM GERIATRIC HEAD INJURY

The topic of geriatric head injury has received more attention in the literature than has any other aspect of geriatric trauma. Unfortunately, all of it is retrospective in nature and, therefore, suffers from many of the same methodological shortcomings discussed above for the remainder of the geriatric trauma literature. These include lack of a specific age definition for geriatric head injury, lack of standardized definitions for specific sub-populations of geriatric head injured patients, and lack of standardized outcome measures. In addition, much of the geriatric head injury literature either provides insufficient details regarding head injury management, or provides results based upon head injury management that would be considered outdated by today's standards. Therefore, it is difficult, and perhaps even dangerous, to make meaningful recommendations regarding the triage of current day geriatric neuro-trauma patients based upon the existing literature. Despite these shortcomings, there is little question that outcomes following traumatic brain injury are much worse in geriatric patients than in their younger counterparts. Vollmer, in a study from the Traumatic Coma Databank, reported on 661 patients age 15 and older with severe brain injuries, defined as GCS less than eight. Mortality for the entire series was 38%, but was 80% for patients greater than 55 years of age. Multi-variate analysis revealed age to be an independent and significant predictor of death and vegetative outcome, beginning at age 45.<sup>15</sup> Another study examined the effect of age upon outcome in patients with acute subdural hematomas. Mortality was 18% in patients between the ages of 18 and 40, but 74% in patients greater than age 65. Once again, advanced age was noted to be a predictive of poor outcomes.<sup>38</sup> In addition to age, a number of other factors have been examined as potential predictors of poor outcome following head injury in geriatric patients. Not surprisingly, the most extensively studied factor is that of admission GCS. Many other factors predictive of poor outcome have been examined including anatomy of the brain injury (epidural versus subdural),<sup>39</sup> need for craniotomy,<sup>15, 40-43</sup> subdural hematoma volume,<sup>38</sup> midline shift,<sup>38,43</sup> pupillary status,<sup>40, 42, 44</sup> and intracranial pressure.<sup>15,45</sup> None of these factors has been examined in sufficient detail to allow us to make any recommendations regarding their potential role as triage tools in geriatric head injury. Therefore, they will not be considered further within this document.

"Low" admission GCS is clearly associated with poor outcomes in elderly head-injured patients. Reuter documented a mortality rate of 87% in elderly patients (age  $\geq 60$ ) with traumatic intracranial hemorrhage and an admission GCS  $< 8$ , though no details regarding head injury management were provided.<sup>46</sup> The available scientific literature, however, does not support the use of a specific GCS that will reliably identify patients destined for poor outcomes. Zietlow, in a study of patients age 65 and older with multi-system injury, identified a GCS  $\geq 8$  as being predictive, while Van Aalst, in a similar study, found a GCS  $\geq 7$  to be associated with death or depending living status.<sup>5, 29</sup> Published studies limited to geriatric patients with head injuries likewise yield no consensus. Rozelle found a GCS  $\geq 7$  to be predictive of hospital mortality in patients with subdural hematomas, and Kilaru noted that this same GCS was associated with a universally poor long-term outcome.<sup>43, 44</sup> Cagetti, however, found that a GCS  $\geq 11$  was associated with a 100% mortality, although this study involved patients 80 years of age and older.<sup>47</sup> Amacher, however, in a similar study of head-injured patients 80 years of age and

older, did achieve an “excellent/good” outcome in a single patient with an admission GCS score in the 3 to 6 range.<sup>48</sup> Thus, it seems that, while “low” GCS scores are indeed associated with poor outcomes, it does not seem possible, or advisable, based upon the existing literature, to make triage decisions in head-injured geriatric patients based solely upon the *admission* GCS. It does seem reasonable to conclude that head-injured patients, 65 years and older, have very poor outcomes when the admission GCS is  $\leq 7$  or 8.

Other authors have examined the prognostic value of the “delayed” GCS score – that is, the GCS score determined 24 hours or more following injury. Both Pennings and Kotwica have advocated a limited course of aggressive therapy in geriatric trauma patients with severe head injuries, although their GCS definitions of futility differ greatly. Kotwica, in a study of head-injured patients 70 years of age and older, noted a 90% mortality in patients with a GCS  $< 9$  when craniotomy was required, and 76% when craniotomy was not required. Based on this finding in 136 patients, they recommend aggressive treatment for 24 hours only for those patients *without* space occupying lesions. Aggressive treatment, then, is continued only in those patients who show “significant” improvement within this time frame.<sup>41</sup> Pennings, in his study of 42 patients age 60 and greater with GCS  $\leq 5$ , concluded that these patients have an extremely poor prognosis, and that if they have not regained “substantial” neurologic function within 24 hours, they are unlikely to do so.<sup>42</sup> Similarly, Ross reported a 100% six month mortality among patients 65 years of age and older who had a persistent GCS  $\leq 8$  at 72 hours following admission.<sup>45</sup> Even though the overall prognosis from geriatric head injury may have improved since these publications due to improvements in head injury management, it is reasonable to expect that these new therapies will exert their maximum effect in the early stages following injury. Thus, in geriatric head injury, it seems reasonable to adopt an initial course of aggressive treatment (with the possible exception of the patient who is moribund upon arrival), followed by a re-evaluation of the patient’s neurologic status at 72 hours post admission. The intensity of the subsequent care provided can then be based upon the initial response to therapy.

## V. SUMMARY

While multiple clinical and demographic factors have demonstrated an association with outcome following trauma in geriatric patients, the ability of any specific factor alone, or in combination with other factors, to predict an unacceptable outcome for any individual geriatric trauma patient is quite limited. An initial course of aggressive therapy (see following section entitled “Geriatric Resuscitation”) seems warranted in all geriatric trauma patients, regardless of age or injury severity, with the possible exception of those patients who arrive in a moribund condition. Geriatric trauma patients who do not respond to aggressive resuscitative efforts within a timely fashion are likely to have poor outcomes even with continued aggressive treatment. Modification of the intensity of treatment provided to these “non-responders” should be considered. For those geriatric trauma patients who do respond favorably to aggressive resuscitative efforts, the prognosis, not only for survival but also for return to their pre-injury level of function, is quite good, and certainly justifies the effort.

## VI. FUTURE INVESTIGATIONS

There are no Class I data that address triage issues in geriatric trauma. Prospective randomized controlled trials are desperately needed which address the prognostic values of age,

injury severity and injury physiology upon ultimate outcome following geriatric trauma. Prior to conducting these studies, there must be agreement concerning the specific age definitions to be used for geriatric trauma, the outcomes to be measured, and the specific clinical criteria that will be used to define pre-existing medical conditions. Furthermore, data generated in such a fashion should be subjected to rigorous and appropriate statistical analysis. Only when a substantial body of literature exists which meets these criteria, will trauma practitioners succeed in providing an appropriate level of care to the geriatric trauma patient based on that patient's predicted outcome.

## VII. REFERENCES

1. Committee on Trauma, American College of Surgeons: *Resources for optimal care of the injured patient: 1999*, Chicago, American College of Surgeons; 1998:14.
2. Champion HR, Copes WS, Buyer D, Flanagan ME, Bain L, Sacco WJ: Major trauma in geriatric patients. *Am J Public Health* 1989;79:1278-1282.
3. McCoy GF, Johnston RA, Duthie RB: Injury to the elderly in road traffic accidents. *J Trauma* 1989;29:494-497.
4. Morris JAJ, MacKenzie EJ, Damiano AM, Bass SM: Mortality in trauma patients: the interaction between host factors and severity. *J Trauma* 1990;30:1476-1482.
5. van Aalst JA, Morris JAJ, Yates HK, Miller RS, Bass SM: Severely injured geriatric patients return to independent living: a study of factors influencing function and independence. *J Trauma* 1991;31:1096-1101.
6. DeMaria EJ, Kenney PR, Merriam MA, Casanova LA, Gann DS: Survival after trauma in geriatric patients. *Ann Surg* 1987;206:738-743.
7. Davis JW, Kaups KL: Base deficit in the elderly: a marker of severe injury and death. *J Trauma* 1998;45:873-877.
8. Gubler KD, Davis R, Koepsell T, Soderberg R, Maier RV, Rivara FP: Long-term survival of elderly trauma patients. *Arch Surg* 1997;132:1010-1014.
9. Osler T, Hales K, Baack B, et al: Trauma in the elderly. *Am J Surg* 1988;156:537-543.
10. Finelli FC, Jonsson J, Champion HR, Morelli S, Fouty WJ: A case control study for major trauma in geriatric patients. *J Trauma* 1989;29:541-548.
11. Smith JSJ, Martin LF, Young WW, Macioce DP: Do trauma centers improve outcome over non-trauma centers: the evaluation of regional trauma care using discharge abstract data and patient management categories. *J Trauma* 1990;30:1533-1538.
12. Ma MH, MacKenzie EJ, Alcorta R, Kelen GD: Compliance with prehospital triage protocols for major trauma patients. *J Trauma* 1999;46:168-175.
13. Phillips S, Rond PC, Kelly SM, Swartz PD: The failure of triage criteria to identify geriatric patients with trauma: results from the Florida Trauma Triage Study. *J Trauma* 1996;40:278-283.
14. Zimmer-Gembeck MJ, Southard PA, Hedges JR, et al: Triage in an established trauma system. *J Trauma* 1995;39:922-928.
15. Vollmer DG: Age and outcome following traumatic coma: why do older patients fare worse? *J Neurosurg* 1991;75:S37-S49
16. Pellicane JV, Byrne K, DeMaria EJ: Preventable complications and death from multiple organ failure among geriatric trauma victims. *J Trauma* 1992;33:440-444.
17. Perdue PW, Watts DD, Kaufmann CR, Trask AL: Differences in mortality between elderly and younger adult trauma patients: geriatric status increases risk of delayed death. *J Trauma* 1998;45:805-810.
18. Knudson MM, Lieberman J, Morris JAJ, Cushing BM, Stubbs HA: Mortality factors in geriatric blunt trauma patients. *Arch Surg* 1994;129:448-453.
19. Horst HM, Obeid FN, Sorensen VJ, Bivins BA: Factors influencing survival of elderly trauma patients. *Crit Care Med* 1986;14:681-684.
20. Shabot MM, Johnson CL: Outcome from critical care in the "oldest old" trauma patients. *J Trauma* 1995;39:254-259.

21. van der Sluis CK, Timmer HW, Eisma WH, ten Duis HJ: Outcome in elderly injured patients: injury severity versus host factors. *Injury*. 1997;28:588-592.
22. Oreskovich MR, Howard JD, Copass MK, Carrico CJ: Geriatric trauma: injury patterns and outcome. *J Trauma* 1984;24:565-572.
23. Broos PL, Stappaerts KH, Rommens PM, Louette LK, Gruwez JA: Polytrauma in patients of 65 and over. Injury patterns and outcome. *Int Surg* 1988;73:119-122.
24. Broos PL, D'Hoore A, Vanderschot P, Rommens PM, Stappaerts KH: Multiple trauma in elderly patients. Factors influencing outcome: importance of aggressive care. *Injury*. 1993;24:365-368.
25. Tornetta P, Mostafavi H, Riina J, et al: Morbidity and mortality in elderly trauma patients. *J Trauma* 1999;46:702-706.
26. Johnson CL, Margulies DR, Kearney TJ, Hiatt JR, Shabot MM: Trauma in the elderly: an analysis of outcomes based on age. *Am Surg* 1994;60:899-902.
27. Battistella FD, Din AM, Perez L: Trauma patients 75 years and older: long-term follow-up results justify aggressive management. *J Trauma* 1998;44:618-623.
28. Carrillo EH, Richardson JD, Malias MA, Cryer HM, Miller FB: Long term outcome of blunt trauma care in the elderly. *Surg Gynecol Obstet* 1993;176:559-564.
29. Zietlow SP, Capizzi PJ, Bannon MP, Farnell MB: Multisystem geriatric trauma. *J Trauma* 1994;37:985-988.
30. DeMaria EJ, Kenney PR, Merriam MA, Casanova LA, Gann DS: Aggressive trauma care benefits the elderly. *J Trauma* 1987;27:1200-1206.
31. Day RJ, Vinen J, Hewitt-Falls E: Major trauma outcomes in the elderly. *Med J Aust*. 1994;160:675-678.
32. Morris JAJ, MacKenzie EJ, Edelstein SL: The effect of preexisting conditions on mortality in trauma patients. *JAMA* 1990;263:1942-1946.
33. Milzman DP, Boulanger BR, Rodriguez A, Soderstrom CA, Mitchell KA, Magnant CM: Pre-existing disease in trauma patients: a predictor of fate independent of age and injury severity score. *J Trauma* 1992;32:236-243.
34. Charlson ME, Pompei P, Ales KL, MacKenzie CR: A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373-383.
35. van der Sluis CK, Klasen HJ, Eisma WH, ten Duis HJ: Major trauma in young and old: what is the difference? *J Trauma* 1996;40:78-82.
36. Shapiro MB, Dechert RE, Colwell C, Bartlett RH, Rodriguez JL: Geriatric trauma: aggressive intensive care unit management is justified. *Am Surg* 1994;60:695-698.
37. Smith DP, Enderson BL, Maull KI: Trauma in the elderly: determinants of outcome. *South. Med J* 1990;83:171-177.
38. Howard MA, Gross AS, Dacey RGJ, Winn HR: Acute subdural hematomas: an age-dependent clinical entity [see comments]. *J Neurosurg* 1989;71:858-863.
39. Rakier A, Guilburd JN, Soustiel JF, Zaaroor M, Feinsod M: Head injuries in the elderly. *Brain Inj*. 1995;9:187-193.
40. Jamjoom A, Nelson R, Stranjalis G, et al: Outcome following surgical evacuation of traumatic intracranial haematomas in the elderly. *Br J Neurosurg*. 1992;6:27-32.
41. Kotwica Z, Jakubowski JK: Acute head injuries in the elderly. An analysis of 136 consecutive patients. *Acta Neurochir.(Wien.)* 1992;118:98-102.

42. Pennings JL, Bachulis BL, Simons CT, Slazinski T: Survival after severe brain injury in the aged. *Arch Surg* 1993;128:787-793.
43. Rozzelle CJ, Wofford JL, Branch CL: Predictors of hospital mortality in older patients with subdural hematoma. *J Am Geriatr.Soc* 1995;43:240-244.
44. Kilaru S, Garb J, Emhoff T, et al: Long-term functional status and mortality of elderly patients with severe closed head injuries. *J Trauma* 1996;41:957-963.
45. Ross AM, Pitts LH, Kobayashi S: Prognosticators of outcome after major head injury in the elderly. *J Neurosci.Nurs.* 1992;24:88-93.
46. Reuter F: Traumatic intracranial hemorrhages in elderly people. *Advances in Neurosurgery* 1989;17:43-48.
47. Cagetti B, Cossu M, Pau A, Rivano C, Viale G: The outcome from acute subdural and epidural intracranial haematomas in very elderly patients. *Br J Neurosurg* 1992;6:227-231.
48. Amacher AL, Bybee DE: Toleration of head injury by the elderly. *Neurosurgery* 1987;20:954-958.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Gubler KD	1997	Long-term survival of elderly trauma patients. <i>Arch Surg</i> ; 132:1010-14.	II	9424	≥67	All (HCFA data base)	4.1% (in-hospital)	Determined 5-year survival of injured cohort compared to 4:1 uninjured cohort matched for age and gender. <i>Injured cohort had more PEC's.</i> Five-year risk of death in injured cohort was 1.7 times that of uninjured cohort and was related to age, gender, ISS and PEC's. Adverse effect of trauma on survival remains long after traumatic episode.
Orskovich MR	1984	Geriatric trauma: injury patterns and outcome. <i>J Trauma</i> ; 24:565-72.	III	100	≥ 70	"severe" injury; [mean ISS = 19]	14% @ 1 month; 15% @ 1 yr	Factors affecting survival included "serious" CNS injury, shock (BP < 80), and burn mechanism. Survival was NOT affected by age, ISS, gender or presence of PEC's. Profile of the non-survivor: required pre-hospital intubation, was in shock at some time, was intubated > 5 days, and developed pulmonary sepsis. Less than 8% of patients were independent at one-year follow-up. Vague definitions of PEC's and entry criteria into study.
Horst HM	1986	Factors influencing survival of elderly trauma patients. <i>Crit Care Med</i> ; 14:681-4.	III	39	>60	ICU admits w/ PAC & Arterial Catheter	31%	Survival related to sepsis and the number of failed organ systems, but NOT age, ISS, TS, APS (Acute Physiology Score), injury mechanism, PEC, presence of shock at admission, or initial cardiopulmonary variables. Small number of patients in this study raises questions regarding the validity of its conclusions.
Amacher AL	1987	Toleration of head injury by the elderly. <i>Neurosurgery</i> ; 20:954-8.	III	56	≥ 80	All head-injury admissions	25%	Overall mortality 25%, but CNS-related mortality was 16%. Seven of eight patients with admission GCS of 3-6 died (87.5% mortality), but the single survivor had an excellent/good outcome. Conversely, 5 of 42 (12%) of patients with GCS of 13 or more died.
DeMaria EJ	1987	Survival after trauma in geriatric patients. <i>Ann Surg</i> ; 206:738-43.	III	82	>65	Excluded burns, penetrating, and isolated ortho injury [ISS=17.9]	21%	Non-survivors were older, had higher ISS's and AIS's for the head and neck, and had more complications. Developed formula to predict outcome based on age, ISS and the presence/absence of cardiac and septic complications. Prospectively tested formula on 61 pts with 92% accuracy. Authors counsel against use of formula (Geriatric Trauma Survival Score) to predict mortality or to limit resuscitative efforts.
DeMaria EJ	1987	Aggressive trauma care benefits the elderly. <i>J Trauma</i> ; 27:1200-06.	III	63	> 65	Excluded burns, penetrating, and isolated ortho injury	Study of survivors only	Examines factors related to home vs nursing home (NH) disposition in elderly trauma survivors. NH pts were older, had higher ISS, more complications, longer LOS's, more severe head and neck trauma, and required surgery more frequently after trauma. These factors are probably not useful for triage purposes as the majority of NH pts ultimately returned home. Overall, 89 % of pts returned home and 57% returned to independent living.
Broos PL	1988	Polytrauma in patients of 65 and over. Injury patterns and outcome. <i>Int Surg</i> 1988; 73:119-22.	III	49	≥ 65	Excluded DOA's and pts who expired prior to any intervention	18% [6 mos] ***** [mean ISS = 33.2]	Factors predicting mortality included coma (not defined) and "early and continued intubation" (intubated pre-hospital or at admission and continued for 5 days or more). Age, ISS and PEC's were not significantly predictive of mortality. Small number of patients on which to make any recommendations. Vague definition of "polytrauma". 76% of survivors returned home.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Osler T	1988	Trauma in the elderly. <i>Am J Surg</i> ; 156:537-43.	III	100	≥ 65	Excluded if pt expired before O.R. or ICU	17%	Factors distinguishing elderly survivors from non-survivors included TS, GCS, ISS, shock, pulmonary sepsis and prolonged ventilation (> 5 days), but not age. Using logistic regression analysis, shock and GCS were found to be the best predictors of geriatric trauma death. No elderly pt survived a TS < 9.
Finelli FC	1989	A case control study for major trauma in geriatric patients. <i>J Trauma</i> ; 29: 541-8.	III	3669	≥ 65	All (MTOS)	18.3%	MTOS data reveals increased trauma mortality beginning at age 45. In the Washington Hospital Center dataset, overall mortality in pts ≥ 65 was twice that of younger pts. ISS-adjusted mortality was greater in the elderly at all ISS levels. ISS was much higher in elderly non-survivors than survivors. Older pts also had higher complication rates. Although no predictive factors for elderly mortality were given, authors recommend triaging elderly trauma victims to trauma centers at a much lower threshold.
Howard MA	1989	Acute subdural hematomas: an age-dependent clinical entity. <i>J Neurosurg</i> ; 71:858-63.	III	67	>65	All patients with acute subdural hematoma > 0.5 cm	74%	Mortality rate of 74% compared to 18% in patients aged 18-40 years. Older patients died significantly later than younger patients (11.2 days vs 2.0 days). Admission GCS was similar for the two groups, but elderly patients had larger subdural hematoma (SDH) volume and more midline shift than younger patients. Advanced age, large SDH volume and midline shift were each predictive of poor outcome, although not independently. The effect, if any, of PEC's on outcome was not studied.
McCoy GF	1989	Injury to the elderly in road traffic accidents. <i>J Trauma</i> ; 29:494-7.	III	312	> 65	All traffic incidents	9.3%	Overall higher mortality in elderly group, even after correcting for ISS. AIS much better predictor of mortality if 1 point is added to the MAIS for patients > 65 years. All pts > 65 yrs w/ MAIS ≥5 died. Small number of patients.
Reuter F	1989	Traumatic intracranial hemorrhages in elderly people. <i>Advances in Neurosurgery</i> ; 17:43-8.	III	64	≥ 60	Included only patients requiring surgery	76%	Mortality was 87% in patients with admission GCS < 8. Mortality also affected by complications with 90% mortality in patients with complications. Description of head injury management not provided.
Morris JA	1990	Mortality in trauma patients: the interaction between host factors and severity. <i>J Trauma</i> ; 30:1476-82.	III	199, 737	≥ 15	All trauma discharges excluding transfers	1.9%	Mortality from minor injury (ISS < 9) increases at age > 65, while for moderate injuries (ISS :9-24), mortality begins to increase at 45 years. ISS is best predictor of mortality in trauma patients, but age, gender and PEC's are also important <i>independent</i> predictive factors of mortality.
Morris JA	1990	The effect of preexisting conditions on mortality in trauma patients. <i>JAMA</i> ; 263:1942-6.	III	3074	≥ 15	All hospitalized trauma deaths in California in 1983	N/A	Case-control study with 4:1 match (survivors:deaths). Trauma mortality increases with increasing numbers of PEC's. PEC's contributing significantly to mortality included liver disease, congenital coagulopathy, COPD, ischemic heart disease and diabetes. . The effect of PEC on mortality was greater in patients w/ ISS < 13, and in pts ≤ 65 years.
Smith DP	1990	Trauma in the elderly: determinants of outcome.	III	456	≥ 65	All patients with	8.6% [mean	Factors associated with outcome included mechanism of injury (burns > MVC > pedestrian struck > assault > falls), and number of

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Smith JS	1990	Do trauma centers improve outcome over non-trauma centers: the evaluation of regional trauma care using discharge abstract data and patient management categories. <i>J Trauma</i> ; 30:1533-8.	III	1332	N/A	All patients with femoral shaft fractures	ISS = 10.8] 1.0% @ trauma centers vs 2.2% @ non-trauma centers	complications. The presence of PEC's was not associated with adverse outcomes, but definition of "PEC" was vague. Low mortality series. PEC's may not influence outcome when ISS and mortality rates are low. Compares outcomes in trauma centers vs non-trauma centers for patients with femoral shaft fractures. Trauma center patients had significantly fewer overall complications (21% vs 33%), and lower mortality. In the subset of patients > 55 years of age, complication rates were 35% at trauma centers, and 47% at non-trauma centers. Elderly trauma patients (age > 55) with significant injuries in addition to their femur fractures were much less likely to be triaged to trauma centers than their younger counterparts (38% vs 70%).
Van Aalst JA	1991	Severely injured geriatric patients return to independent living: a study of factors influencing function and independence. <i>J Trauma</i> ; 31:1096-1101.	III	98	≥ 65	Blunt trauma; ISS ≥ 16	44% in hospital	1 to 6 year follow-up (mean 2.82 years) of 54 elderly blunt trauma patients with ISS ≥ 16, who survived initial hospitalization. 11% of these died during the follow-up period, and only 17% regained their pre-injury function. 67% however returned to independent living. Factors associated with a poor outcome (death or dependent living status) included GCS ≤ 7, age ≥ 75, shock on admission, presence of head injury (AIS-Head ≥ 3), and sepsis.
Vollmer, DG	1991	Age and outcome following traumatic coma: why do older patients fare worse? <i>J Neurosurg</i> ; 75:S37-49.	III	661	≥ 15	Traumatic Coma Data Bank (TCDB) patients; age ≥ 15, GCS ≤ 8, gunshot wounds to head and patients meeting brain death criteria on arrival were excluded]	38%	Reports outcomes at 6 months post injury for patients with severe brain injuries (GCS < 8). Overall mortality was 38%, but was 80% for patients > 55 years of age. No "elderly" patient made a "good" recovery, and there were fewer "elderly" patients with moderate disability, severe disability and vegetative survival compared to younger patients. Early (<48 hours) mortality was similar among all age groups, but late (>48 hours) mortality was significantly higher in older patients. Although pre-existing medical conditions and complications were more frequent in elderly patients and, thus, were associated with poor outcome, multivariate analysis revealed age to be an <i>independent</i> and significant predictor of death and vegetative outcome, beginning at age 45. Whether pre-existing medical conditions and complications remain <i>as independent</i> predictors of poor outcome is not stated. The authors conclude that the poor outcome following head injury in "elderly" patients is primarily due to the limited capacity of the aging brain to recover following injury.
Cagetti B	1992	The outcome from acute subdural and epidural intracranial haematomas in very elderly patients. <i>Br J Neurosurg</i> ; 6:227-32.	III	28	≥ 80	Excluded patients with intracerebral hematomas and con-	88%, (compared to 57% in pts	All patients with GCS ≤ 11 died. Pre-existing diseases and multiple systems organ failure accounted for the majority of deaths. All surviving patients successfully returned to their pre-injury state of health. No significant difference between the volume of clot or the frequency of associated cerebral contusions between those patients ≥ or < 80 years.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Janjoom A	1992	Outcome following surgical evacuation of traumatic intracranial haematomas in the elderly. <i>Br J Neurosurg</i> ; 6:27-32.	III	66	≥65	All patients undergoing craniotomy for evacuation of post-traumatic hematoma	61%	The authors conclude the level of consciousness at the time of operation correlates with outcome better than do other parameters. No description of management was provided, making it difficult to determine whether care provided to the two populations was equivalent.  Increased mortality (86%) in the subset of patients ≥ 80 years of age. Outcome also worse if craniotomy performed within 24 hours of injury. Authors feel craniotomy not justified in patients with a pre-operative GCS of 4 or less or in those with unilateral or bilateral pupillary dilatation since all patients in these 2 categories had poor outcomes (Glasgow Outcome Score of 1-3).
Kotwica Z	1992	Acute head injuries in the elderly. An analysis of 136 consecutive patients. <i>Acta Neurochir (Wien)</i> ; 118:98-102.	III	136	> 70	Head Injury Only	52%	For pts with GCS < 9, mortality was 90% when craniotomy was required, and 76% when craniotomy was not required. Based on these results, authors recommend limiting therapy in pts with GCS < 9 with space-occupying lesions. In patients without space-occupying lesions and GCS < 9, authors recommend aggressive treatment for 24 hours, and limiting further treatment to those with significant improvement by this time. No statistical analysis performed. Small number of patients on which to base such critical recommendations.
Milzman DP	1992	Pre-existing disease in trauma patients: a predictor of fate independent of age and injury severity score. <i>J Trauma</i> ; 32:236-43.	III	7798	≥ 15	All admissions w/ ISS > 1; Excluded if survival < 24 hours or cardiac arrest on arrival	9.2% (PEC+) _____ 3.2% (PEC-)	4 year retrospective study at a single Level I trauma center. Trauma mortality increases with increasing numbers of PEC's. The effect of PEC on mortality is independent of age and ISS, but becomes less important at age > 55 years or at ISS > 20.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Pellicane JV	1992	Preventable complications and death from multiple organ failure among geriatric trauma victims. <i>J Trauma</i> ; 33:440-4.	III	374	≥ 65	Burns Excluded	8%	Elderly non-survivors were significantly older, had higher ISS and lower TS than elderly survivors. TS < 15 was associated with a 45 % mortality, but 52 % of deaths occurred in pts w/ a TS ≥ 15. Potentially preventable complications contributed to mortality in 62 % of organ failure deaths, and one third of sudden deaths. 70% of organ failure deaths in the TS 15-16 group were contributed to by potentially preventable complications.
Ross AM	1992	Prognosticators of outcome after major head injury in the elderly. <i>J Neurosci Nurs</i> ; 24:88-93.	III	195	≥ 65	GCS ≤ 8 or intracranial hematoma requiring evacuation	20% @ 72 hours ----- 75% @ 6 months	In patients w/ admission GCS ≤ 8, 83% were still in coma after 72 hours. All of these patients died within 6 months. Patients with ICP's ≥ 20 had higher 72 hour and 6 month mortality, and greater 72 hour neurologic disability compared to patients with ICP's < 20. However, incidence of shock and apnea were greater in elevated ICP group, which could have adversely affected neurologic outcome and mortality. Study describes patients treated between 1978 and 1988. Conclusions might therefore have limited applicability to current patient care.
Broos PL	1993	Multiple trauma in elderly patients. Factors influencing outcome: importance of aggressive care. <i>Injury</i> ; 24:365-8.	III	126	≥ 65	Excluded DOA's and pts who expired prior to any intervention	17% [6 mos] ***** [mean ISS = 33.2]	Factors predicting mortality included GCS ≤ 8 and "early and continued intubation" (intubated pre-hospital or at admission and continued for 5 days or more). Age, ISS and PEC's were not significantly predictive of mortality. Vague definition of "multiple trauma". Within 6 months of discharge, 78% of patients had returned to their pre-injury surroundings.
Carrillo EH	1993	Long term outcome of blunt trauma care in the elderly. <i>Surg Gynecol Obstet</i> ; 176:559-64.	III	94	≥ 65	Excluded burns, penetrating, isolated ortho injury, and pts with minimal injuries.	13%	Mortality correlated well with APACHE II, but combination of APACHE II and ISS performed better than APACHE II alone. All patients with APACHE II ≥ 15 and ISS ≥ 30 died, but this accounted for only 1/3 of all deaths in the series. At 1 to 3 year follow-up, 84 % of patients surviving hospital discharge were independent at home.
Penning's JL	1993	Survival after severe brain injury in the aged. <i>Arch Surg</i> ; 128:787-93.	III	42	≥ 60	Excluded if GCS > 5, penetrating injury, pts w/ nl CT, pts expiring w/i 6 hours	79%	Of 9 survivors, 6 were in a persistently vegetative state, and 2 were severely disabled. The final survivor was moderately disabled and was discharged home. After discharge, older pts tended to deteriorate neurologically, while younger pts tended to improve or remain stable. Factors predictive of mortality were a decreased 6 hour GCS, age ≥ 60, lack of need for craniotomy, cerebral edema, and non-reactive pupils. Authors conclude that pts > 60 years w/ GCS < 5 have an extremely poor prognosis, and that if they do not regain substantial neurologic function within 24 hours, they are unlikely to do so.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Day RJ	1994	Major trauma outcomes in the elderly. <i>Med J Aust</i> ; 160:675-8.	III	118	>60	ISS > 15	30.5% (early); 31% (late)	Mean ISS =25. Minimum 2 year (average 3 year) follow-up obtained. Late mortality much higher in patients > 70 years old (50%) than in patients 61-70 years old (8%). Of survivors, 81% were living independently, and 76% scored maximally on ADL testing. Authors claim "that age is a significant factor in long term survival after major trauma", but no supporting statistical analysis provided.
Johnson CL	1994	Trauma in the elderly: an analysis of outcomes based on age. <i>Am Surg</i> ; 60:899-902.	III	289	≥ 65	SICU admits only	16.3%	Despite similar mean ISS, elderly pts had higher admission SAPS (Simplified Acute Physiology Score) compared to younger pts. SICU mortality increased with increasing ISS and SAPS, though ISS-adjusted mortality not statistically different between elderly and younger pts. Authors conclude that injury <i>physiology</i> (SAPS) better predictor of early death, while age still important predictor of death after ICU discharge.
Knudson MM	1994	Mortality factors in geriatric blunt trauma patients. <i>Arch Surg</i> ; 129:448-53.	III	852	≥ 65	Blunt Trauma Only	18.4%	Factors predictive of <i>in-hospital</i> mortality were: male gender; injury mechanism; ISS; injuries to brain, chest or abdomen; TS; and RTS. Admission physiologic factors associated with death included: BP < 90; RR < 10, TS, <7; and a GCS = 3. A TS < 7 was associated with 100 % mortality, as was a RR < 10. ISS was best predictor of mortality, although ISS not "available" as a prognostic variable at admission.
Shapiro MB	1994	Geriatric trauma: aggressive intensive care unit management is justified. <i>Am Surg</i> ; 60:695-98.	III	170	≥ 60	All trauma admissions	21.8%	All deaths were in ICU patients. ICU mortality correlated with the number of organ systems failing and with severe head injury (not defined). Survival not related to the presence of PEC's.
Zietlow SP	1994	Multisystem geriatric trauma. <i>J Trauma</i> ; 37:985-8.	III	94	≥ 65	ISS ≥ 10 [mean ISS = 18]	23%, in-hospital	Factors predictive of death (univariate): severe brain injury (GCS ≤ 8), inotropic/ventilatory support, previous MI, shock, chronic renal insufficiency and bradycardia. Factors predictive of death (multivariate): severe brain injury (GCS ≤ 8) and previous MI. At mean follow-up of 12 months, 75% of pts were at home and independent and 49% were back to their normal level of activity.
Rakier A	1995	Head injuries in the elderly. <i>Brain Inj</i> ; 9:187-93.	III	263	≥ 65	Consecutive series of head injuries, including concussions	17.5%	High mortality rates noted in patients with cerebral contusions (~28% mortality), and acute subdural hematomas (33% mortality). All patients with acute epidural hematomas had poor outcomes. Overall conclusions weakened by lack of data on admission GCS, author's grouping of patients according to predominant finding on head CT (only one finding allowed per patient), and lack of long-term follow-up.
Rozzelle CJ	1995	Predictors of hospital mortality in older patients with subdural hematoma. <i>J Am Geriatr Soc</i> ;	III	157	≥ 65	Pts w/ traumatic subdural hematomas	30.6%	Factors predictive of hospital mortality included GCS ≤ 7, age ≥ 80, <i>acute</i> duration of hematoma, and need for craniotomy. Presence of co-morbidities, use of antithrombotics, and midline shift on CT did not influence outcome.

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Shabot MM	1995	Outcome from critical care in the "oldest old" trauma patients. <i>J Trauma</i> ; 39:254-9.	III	45	? 75	All trauma admissions to the SICU	28.9%	"Oldest" patients had significantly higher ISS values. Although TS was similar between all age groups, SAPS (both on the first SICU day and also maximum SICU SAPS) increased significantly with age. In general, increases in ISS in the "oldest" group translated into larger increases in SAPS, indicating that these patients have a progressively greater physiologic response to a given level of injury. Factors predictive of overall SICU mortality (all ages) included ISS, TS, SAPS, and day of maximum SAPS. Age itself not predictive of overall SICU mortality, and when stratified by SAPS, mortality in "oldest" patients is similar to younger patients.
Zimmer-Gembeck MJ	1995	Triage in an established trauma system. <i>J Trauma</i> ; 39:922-8.	III	26025	N/A	ISS $\geq$ 1	N/A	Examines success of pre-hospital trauma triage. Assumes all pts w/ ISS between 1 and 9 should have been triaged to a non-trauma hospital, while all pts with ISS $\geq$ 16 should have been triaged to a trauma center. Undertriage rate for entire study population was 21%, but was 56 % for patients > 65 years of age. Overtriage of elderly trauma patients was only 10% (28% for entire study population). In addition to the problem of undertriage in the elderly trauma patient, this study also found that most trauma deaths in non-trauma hospitals were in elderly patients with ISS between 1 and 9.
Kilaru S	1996	Long-term functional status and mortality of elderly patients with severe closed head injuries. <i>J Trauma</i> ; 41:957-63.	III	40	> 65	GCS $\leq$ 8; DOA's and inaccurate GCS's were excluded	68%	Overall mortality at average 38 month follow-up was 73%. Factors predictive of mortality were GCS, GCS-motor response, and fixed pupils. ISS was not found to be predictive of mortality, but age and TS showed a trend towards significance. With multiple regression analysis, only GCS and heart rate correlated with death. All pts w/ admitting GCS = 3 died in hospital, and all with GCS $\leq$ 7 either died, were vegetative, or had severe disabilities. On long-term follow-up, neurologic function improved very little after hospital discharge. Overall weak conclusions due to small number of patients and limited use of ICP monitoring.
Phillips S	1996	The failure of triage criteria to identify geriatric patients with trauma: results from the Florida Trauma Triage Study. <i>J Trauma</i> ; 40:278-83.	III	3980	$\geq$ 55	Excluded if burns, isolated hip fx, interhospital transfers, and incomplete data	2.4 %	Study uses AIS data and several assumptions to determine pt's "ideal" triage destination (trauma center vs non-trauma center), and then compares actual pt triage destination with "ideal" destination. Overtriage in the elderly population was 7.4% (compared to 11.3% in younger patients, and target overtriage rate of 20%). Undertriage in the elderly group was 71% (compared to 36% in younger group, and target undertriage rate of 5%). Triage criteria failed to identify nearly all elderly major trauma cases from falls.
Van der Sluis CK	1996	Major trauma in young and old: what is the difference?	III	121	$\geq$ 60	ISS $\geq$ 16	38.8%	Compares outcome of elderly with that of younger patients, all with ISS $\geq$ 16. Mortality in both groups increased with increasing ISS, and ISS was similar for the 2 groups. ISS in elderly non-survivors was higher

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
Van der Sluis CK	1997	Outcome in elderly injured patients: injury severity versus host factors. <i>Injury</i> ; 28:588-92.	III	42	> 60	ISS $\geq$ 16	31%	Compares outcomes between elderly trauma pts and elderly hip fracture pts. In-hospital mortality was much higher for the trauma elderly (31%) than for the hip fracture elderly (3%), but long-term survival (7-8 years post-trauma) was similar (29%). Higher late mortality in hip fracture group ascribed to higher incidence of poor "pre-injury medical status" (very loosely defined) in this pt population (53% vs 12%). Predictors of late mortality included age, poor pre-injury medical status, and male gender, while early mortality is more a function of ISS.
Battistella FD	1998	Trauma patients 75 years and older: long-term follow-up results justify aggressive management. <i>J Trauma</i> ; 44:618-23.	III	279	$\geq$ 75	All Trauma Service admissions. Mean ISS = 9.4	23%, including DOA's	4 year follow-up obtained on 81 % of the 279 patients who survived to hospital discharge. Poor outcome (survival < 6 months following discharge) predicted by pre-existing dementia, hypertension and COPD, but not by age or ISS. Of long term survivors, 83% living independently.
Davis JW	1998	Base deficit in the elderly: a marker of severe injury and death. <i>J Trauma</i> ; 45:873-7.	III	274	$\geq$ 55	Excluded pts w/ initial ABG obtained > 1 hour after injury	varied with base deficit	Correlated admission base deficit (ABD) with mortality, ISS and ICU-LOS. Higher mortality in elderly despite similar ISS and ABD. In pts $\geq$ 55 yrs, an $ABD \leq -6$ was associated with a 67% mortality and a 78% PPV for an ISS $\geq$ 16. However, less severe ABD's (> -6) were still associated with significant mortality (24%). Even a normal ABD (-2 to 2) in this age group was associated with an 18% mortality.
Perdue PW	1998	Differences in mortality between elderly and younger adult trauma patients: geriatric status increases risk of delayed death. <i>J Trauma</i> ;45: 805-10.	III	448	$\geq$ 65	Excluded if isolated single-system injury admitted to non-trauma service	14%	Elderly mortality twice that of younger patients., and was more delayed, with the majority occurring more than 24 hours after admission. Factors predictive of early (< 24 hours) mortality included ISS, RTS, and age. Factors predictive of late (>24 hours) mortality were ISS, RTS, age, pre-existing cardiovascular or liver disease, and the development of cardiac, infectious or renal complications.
Ma MH	1999	Compliance with prehospital triage protocols for major trauma patients. <i>J Trauma</i> ; 46:168-75.	III	32950	$\geq$ 55	All trauma transports	N/A	Documents compliance of pre-hospital providers with pre-hospital trauma triage criteria. Compliance with <i>anatomic</i> criteria was 86%, and did not vary with patient age. Compliance with <i>physiologic</i> and <i>mechanism</i> criteria was poor (34% and 46% respectively), and was statistically worse for pts $\geq$ 55 years of age, compared to younger pts.
Tornetta P	1999	Morbidity and mortality in	III	326	$\geq$ 60	Blunt	18.1%	Factors predictive of mortality included transfusion, ISS (particularly

## Evidentiary Table: Triage Issues in Geriatric Trauma

First Author	Year	Reference	Data Class	# Pts	Age	Pt. Population	Mort.	Conclusion
		elderly trauma patients. <i>J Trauma</i> ; 46:702-6.				trauma only; slip-and-fall injuries were excluded		AIS-Head and Neck and AIS-Thorax), GCS and fluid requirement. In addition, sepsis, ARDS and MI were significant risk factors for mortality. Geriatric Trauma Survival Score was not predictive of survival.

# PARAMETERS FOR RESUSCITATION OF THE GERIATRIC TRAUMA PATIENT

## I. Statement of the problem

There is no doubt that the elder trauma patient presents trauma surgeons with a complex challenge. The effects of aging upon individual organ systems and the presence of comorbid conditions combine to create a milieu that does not allow for errors in resuscitation or delays in diagnosis. It is widely known that geriatric patients have less physiologic reserve than younger patients and that mortality rates are higher than in a younger cohort. There is a growing sentiment that the conduct of resuscitation for the injured elder must be undertaken with an aggressive and thoughtful approach. Outcome data suggests that the elderly benefit by an aggressive approach to resuscitation. It is believed by some that the pulmonary artery catheter should be a routine part of the resuscitation process for the severely injured geriatric patient. In addition, there are certain laboratory assays that have been recommended for use in this clinical scenario. There is confusion, however, regarding end-points for resuscitation and which patients benefit from invasive hemodynamic monitoring.

## II. Process

Literature used for these guidelines was obtained via a search of the MEDLINE database from the National Library of Medicine. Citations in the English language during the period of 1966 through 1999 using the words elderly, geriatric, trauma, shock, and resuscitation were identified. Citations concerned primarily with multisystem trauma or single organ injury in a multisystem context were utilized. Additional non-trauma references were used to relate epidemiological or physiologic factors concerning the geriatric patient to the context of potential injury. This search identified 4,783 references. For use in the evidentiary table, these were then sorted in order to identify articles associated with geriatric trauma patients exclusively. The bibliographies of each article were searched for additional references not identified by the original MEDLINE query. Letters to the editor, case reports, review articles, and series examining non-trauma patients were excluded for use in the evidentiary table. The references were classified by methods employed by the Canadian and U. S. Preventative Task Force. Classification of references was graded based on the strength of the scientific evidence. For purposes of practice management guidelines for trauma, data were classified as follows:

Class I evidence: Prospective randomized controlled trials (PRCTs) - the gold standard of clinical trials. Some may be poorly designed, have inadequate numbers, or suffer from other methodological inadequacies, and thus may not be clinically significant.

Class II evidence: Clinical studies in which the data were collected prospectively, and retrospective analyses which were based on clearly reliable data. These types of studies include observational studies, cohort studies, prevalence studies and case control studies.

Class III evidence: Most studies based on retrospectively collected data. Evidence used in this class includes clinical series, databases or registries, case reviews, case reports, and expert opinion.

### III. Recommendations

#### A. Level I

There are insufficient data to support a level I recommendation for the method and end-points of resuscitation in the elderly patient as a standard of care.

#### B. Level II

1. Any geriatric patient with physiologic compromise, significant injury (AIS > 3), high risk mechanism of injury, uncertain cardiovascular status, or chronic cardiovascular or renal disease should undergo invasive hemodynamic monitoring using a pulmonary artery catheter.

There are insufficient data to support a level II recommendation for the method and end-points of resuscitation in the elderly patient as a standard of care.

#### C. Level III

1. Attempts should be made to optimize to a cardiac index  $\geq 4$  L/min/M<sup>2</sup> and/or an oxygen consumption index of 170 cc/min/M<sup>2</sup>.

2. Base deficit measurements may provide useful information in determining status of resuscitation and risk of mortality.

### IV. Scientific Foundation

It is widely known that the citizenry of the U.S. is continuing to age.<sup>1,2,3</sup> The elderly population (65 years and older) increased 11-fold from 1900 to 1994, while the segment under the age of 65 increased only 3-fold during the same period.<sup>2</sup> Data from the United States government shows that the life expectancy of the U.S. population reached 76.5 years the highest at any time in U.S. history.<sup>1</sup> There will be a dramatic increase in the elderly population due to the aging of the “baby-boom” generation (75 million babies born between 1946 and 1964).<sup>2</sup> While projection assumptions vary, using the Census Bureau’s “Middle Series” projections (moderate fertility, mortality, and immigration assumptions) the elderly will make up 12.8% by 2000 and 20.4% by 2050.<sup>2</sup>

Trauma ranks as the fifth leading cause of death when considering all races, both sexes, and all ages.<sup>1</sup> For years 65 and over, trauma ranks seventh, although the rate per 100,000 is 92.1 compared with 35.7 for all age groups. Unlike younger age groups, there is relatively little variation in death rates between black and white races. These data indicate that in the future there will be an unprecedented number of elderly persons at risk for injury.

Advancing age is associated with a gradual decline in organ function. Problems due solely to senescence and diseases not associated with age may be difficult to distinguish from one another, but it is important to account for all disorders concomitant with the injury. The walls of the heart become less compliant and cardiac index decreases 1% per year with age while systemic vascular resistance rises 1% per year.<sup>4</sup> Maximum heart rate is also reduced with age. In addition, the heart is less able to respond to the stress of injury as there is an age-related decrease in the effectiveness of adrenergic stimulation.<sup>5</sup> The prevalence of hypertension also increases as a function of age. In the United States, 59.2% of white males aged 65-74 are hypertensive and this

increases dramatically to 82.9% in elderly black females.<sup>6</sup> The end result of these age-related changes is a decreased ability to respond to the stress of injury or critical illness.

There are numerous changes in respiratory function with increasing age. The chest wall becomes less compliant and the elasticity of the lung decreases.<sup>4</sup> The loss of compliance results in a greater dependence on diaphragmatic breathing.

Renal mass is rapidly lost after the age of 50 and a corresponding fall in glomerular filtration rate occurs beyond the age of 60 due to the loss of nephrons.<sup>4</sup> Measurement of creatinine clearance becomes more important in the geriatric patient, since serum creatinine may be lowered as a result of decreased muscle mass, giving a false sense of security with respect to renal function. Age-related vascular changes result in a decreased percentage of blood flow to the older kidney.<sup>4</sup>

Endocrine function is also seen with advancing age. The production and turnover of thyroid hormone species is significantly reduced.<sup>7</sup> Tissue responsiveness to thyroid hormone is lessened, resulting in striking similarities between clinical hypothyroidism and the changes commonly seen in the elderly as a result of senescence.<sup>7</sup> Normal adrenal function is critical in order to respond to the stress of injury and critical illness. Basal, circadian, and stimulated cortisol secretion remains intact with aging.<sup>8</sup> There is an age-related decrease in the catabolism of cortisol, although this is compensated with a decrease in the rate of catabolism.

The question of whether or not preexisting disease contributes to poor outcome has not yet been conclusively answered. The prevalence of comorbid conditions in trauma patients is between 8.8% and 19.3%.<sup>9</sup> In injured patients greater than 65 years however, the incidence climbs to 30%.<sup>10</sup> Milzman found that by 75 years, 69% of patients had one or more pre-existing conditions.<sup>11</sup> Smith et al. found at least one comorbidity in 61.6% of patients in their series.<sup>12</sup> In a study of 102 patients from Switzerland admitted with femur fractures, 16% presented with a single comorbid condition, 45% presented with two comorbid conditions, 28% with three conditions, and 11% presented with four.<sup>13</sup> Battistella found an average of two pre-existing medical problems in injured patients 75 years and older.<sup>14</sup> After controlling for age, Sacco found that hepatic, cardiovascular, respiratory, renal, and diabetes adversely affected survival.<sup>15</sup> Milzman and MacKenzie found that mortality and length of stay increased as the number of pre-existing conditions rose.<sup>11, 16</sup>

Criteria for hemodynamic monitoring are not clear in this population. The gravity of this situation is underscored since it has been found that the elderly patient is more likely to present in shock than younger patients with similar trauma and injury severity scores.<sup>17</sup> In geriatric patients undergoing elective surgery, occult physiologic compromise has been shown to contribute to poor outcome. DelGuerico found significant physiologic compromise in geriatric patients who had been "cleared" for elective surgery.<sup>18</sup> Among those who were not able to be optimized prior to surgery, all died postoperatively. Similar work has also been done in trauma patients.

Scalea et al. found significant measurable hemodynamic compromise in elderly patients who were clinically stable after initial evaluation after blunt multiple trauma.<sup>19</sup> Based upon institutional experience, criteria were developed to select patients for invasive hemodynamic monitoring. These criteria included: pedestrian-motor vehicle mechanism, initial blood pressure < 150 mm Hg, acidosis, multiple fractures, and head injury. Patients were moved to the intensive care unit as quickly as possible. Pulmonary artery catheters and arterial lines were inserted in all patients. Volume infusion and ionotropes were used to augment hemodynamic parameters. Attempts were made to optimize patients to a cardiac index  $\geq 4 \text{ L} / \text{min} / \text{M}^2$  or an oxygen

consumption index of 170 cc / min / M<sup>2</sup>. Thirteen of 30 patients were found to be in cardiogenic shock and 54% of these died. There were statistically significant differences between optimized cardiac output and systemic vascular resistance in survivors compared with non-survivors. The vital message from this important work is that a multiply injured geriatric patient may appear “stable” yet have a profound perfusion deficit from a dangerously low cardiac output. The early use of invasive hemodynamic monitoring will identify this deficit and afford the opportunity to help improve survival.

The only randomized data concerning resuscitation in geriatric patients was conducted by Schultz et al.<sup>20</sup> These authors studied the role of physiologic monitoring in patients with fractures of the hip. Seventy patients were randomly divided into a monitored group and a control group. A central venous line was placed into the control group and a pulmonary artery catheter into the monitored group. The mean age for the nonmonitored group was 67 years (range, 40-89) and that for the monitored group was 78 years (range, 40-95). Based upon the data obtained, physiologic abnormalities were “appropriately corrected.” Postoperative morbidity was similar between the two groups. The postoperative mortality in the monitored group was 2.9% and the mortality in the nonmonitored group was 29%. The primary weakness in this study is that no clear parameters are provided to guide resuscitation. This study evaluated patients with hip fractures and not the multisystem elderly trauma patient.

Tornetta et al. retrospectively reviewed 326 patients at four hospitals who were 60 years of age or greater.<sup>21</sup> Using univariate analysis, patients who died displayed significantly greater transfusion requirement (10.9 vs. 2.9 units) and fluid infusion (12.4 vs. 4.9 liters). Both transfusion and fluid requirements were found to be predictive of mortality. The authors concluded that the risks of invasive monitoring are justified in patients with ISS > 18, but for patients with ISS < 18, indications need to be clarified.

The importance of shock and fluid replacement in the elder trauma patient was addressed by Oreskovich et al.<sup>22</sup> One-hundred consecutive elderly patients (mean age 74 years) were followed for a minimum of one year. A profile of the non-survivor was constructed: 1. Pre-hospital intubation (93% mortality), 2. Shock (100%), 3. Intubated greater than 5 days (100%), and 4. Gram-negative pulmonary sepsis (80%). All non-survivors were in shock (systolic blood pressure < 80 mm Hg) for at least 15 minutes between injury and admission. Only 6% of survivors were found to be in shock. During this study the protocol for prehospital care in the hypotensive patient called for 2,200 ml. lactated Ringer’s solution prior to arrival at hospital.

Perdue et al. retrospectively studied 4,691 patients aged 16-64 years and compared these to 448 patients aged 65 years or greater.<sup>23</sup> Elderly mortality was 14% compared to 6% in the younger cohort and the difference was statistically significant (p < 0.001). After controlling for Injury Severity Score, Revised Trauma Score, pre-existing disease, and complications, the elderly were 4.6 times as likely to die compared to the young. The author’s practice is to admit elderly patients to ICU if they have significant injury (AIS > 3), shock, or significant chronic cardiovascular or renal disease. Pulmonary artery catheters were not placed unless volume or cardiac status is uncertain.

Knudson et al. retrospectively analyzed physiologic status in 852 blunt trauma patients aged 65 years or older.<sup>24</sup> Mortality rose with a decreasing Trauma Score and mortality was 100% with a Trauma Score < 7. Each individual component of the TS was found to be predictive of mortality when analyzed independently. Systolic blood pressure < 90 mm Hg associated with 82% mortality rate. Multiple logistic regression used to construct formula to help predict which patients may benefit from aggressive resuscitation.

Physiologic status was also addressed by Pellicane et al.<sup>25</sup> The authors reviewed 374 consecutive trauma patients over the age of 65 years. Trauma score was significantly ( $p < 0.001$ ) higher in patients who survived. Mortality was significantly ( $p < 0.05$ ) increased in patients with  $TS < 12$  (65%) and  $TS = 12-14$  (25%) when compared with patients with  $TS = 15-16$  (5%). The authors concluded that geriatric patients with a  $TS < 15$  are at high risk for complications and should be admitted to the ICU and treated aggressively.

Horst et al. retrospectively studied 39 trauma patients over the age of 60 years.<sup>26</sup> Patients were admitted to the intensive care unit and monitored with arterial and pulmonary artery catheters. Fifteen (38%) of patients presented with shock (systolic blood pressure  $< 80$  mm Hg). Although survivors tended to have higher mean arterial pressure, cardiac index, left ventricular stroke work, and oxygen delivery, the differences compared with non-survivors were not statistically significant.

The importance of shock was further underscored by Van Aalst et al.<sup>27</sup> The authors retrospectively analyzed 98 geriatric ( $> 65$  years) blunt trauma patients with  $ISS \geq 16$ . Of 48 surviving patients, only one patient presented in shock. This is compared with 50 patients who died where 15 presented in shock. The presence of shock (systolic blood pressure  $< 90$  mm Hg) upon admission was the most significant factor associated with a poor outcome. Sepsis was also identified as a factor also contributing to poor outcome.

The state of resuscitation as evaluated by base deficit was evaluated by Davis et al.<sup>28</sup> The authors studied the utility of base deficit in 274 patients 55 years and older. Arterial blood gases obtained within one hour after admission. There was a statistically significant increase in mortality with increasing base deficit. Compared to a younger cohort, mortality in the elderly was significantly increased for a given base deficit despite similar Injury Severity Score. The positive predictive value of base deficit for significant injury was similar between young and old, but the negative predictive value was significantly better in younger patients. The authors concluded that a base deficit  $> 6$  is particularly ominous in elderly.

The above data indicate the use of necessary hemodynamic monitoring and careful trending of vital signs rather than relying on a single set of "normal" vitals.<sup>29</sup> Since the elderly patient is often not able to generate an augmented cardiac output in response to hemorrhage, early invasive hemodynamic monitoring and judicious use of vasoactive drugs (after appropriate fluid resuscitation) as recommended by Scalea should be recommended for any geriatric patient with significant injuries.

#### **IV. Evidentiary Table**

The articles listed in the evidentiary table are those utilized to formulate these guidelines for conduct of resuscitation and the use of invasive hemodynamic monitoring in resuscitation of the geriatric trauma patient. The data are listed in alphabetical order by first author of the publication. Included are: 1 Class I article, 1 Class II article and 7 Class III articles.

#### **V. Summary**

The elderly (65 years and older) are the fastest growing segment of the United States population. While trauma is only the seventh leading cause of death in the elderly, the incidence of injury (per 100,000) is significantly higher when compared to a younger cohort. United States

Bureau of Census data indicate that in the future there will be an unprecedented number of elderly persons at risk for injury.

It is widely known that the elderly display a high incidence of premorbid conditions. However, the question of whether or not preexisting disease contributes to poor outcome after injury has yet to be conclusively answered. Several studies have indicated that shock, respiratory failure, decreasing trauma score, increasing injury severity score, increasing base deficit, and infectious complications portend a poor outcome in the elderly.

Data indicate that the multiply injured geriatric patient may appear “stable” yet have a profound perfusion deficit secondary to low cardiac output. The early use of invasive hemodynamic monitoring may afford the opportunity to help improve survival.

Although the injured elder is more likely to die than the younger patient, an aggressive treatment program will allow many geriatric patients to regain their preinjury independence. Attention to detail, while important for all trauma patients, must be heightened in the injured elder as the opportunity for good outcomes may be fleeting.

## **VI. Future Studies**

The paucity of literature evaluating the conduct and end-points of resuscitation of the geriatric trauma patient requires that further clinical work be conducted. Randomized trials in severely injured geriatric patients must be done in order to determine which patients would benefit from invasive monitoring and the end-points which should be used for completing the resuscitation. Trials like this however have ethical, medicolegal, and methodological implications that may prevent their inception.

There are many parameters that have been shown to correlate with poor outcome in this population. We have no control over some of these, such as patient age. Some can be controlled with prevention techniques, as in the case of the pedestrian-motor vehicle crash, which has been shown to be associated with mortality. As clinical practitioners, we should focus our efforts on those areas where we would be able to exert an impact. The shock state, acidosis, and sepsis have been shown to directly correlate with mortality. Aggressive identification, correction, and monitoring of these pathophysiologic states may be able to improve outcome. Laboratory assays, such as base deficit, may have promise for measuring the adequacy and completeness of resuscitation. Certain drugs such as beta-blockers have been shown to improve outcome in elderly general surgery patients, but have not yet been studied in trauma patients.

## REFERENCES

1. Hoyert DL, Kochanek KD, Murphy SL. Deaths: final data for 1997. National Vital Statistics Reports from the Centers for Disease Control and Prevention. Available at: [http://www.cdc.gov/nchswww/data/nvs47\\_19.pdf](http://www.cdc.gov/nchswww/data/nvs47_19.pdf). Accessed August 13, 1999.
2. U.S. Bureau of the Census. Current Population Reports, Special Studies, P23-190, 65? in the United States. Available at: <http://www.census.gov/prod/1/pop/p23-190/p23-190.html>. Accessed 7/31/99.
3. Markus GR. The graying of America: major Social Security and Medicare battles are just beginning. *Bull Am Coll Surg* 1997; 82:25-30.
4. Fairman R, Rombeau JL. Physiologic problems in the elderly surgical patient. In: Miller TA, Rowlands BJ, eds. *Physiologic Basis of Modern Surgical Care*. St. Louis: The C.V. Mosby Company; 1988:1108-1117.
5. Lakatta EG. Age-related alterations in the cardiovascular response to adrenergic mediated stress. *Fed Proc* 1980; 39:3173-3177.
6. Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure: The 1992 Report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC-V). *Arch Intern Med* 1993; 153:154-183.
7. Mooradian AD. Normal age-related changes in thyroid hormone economy. *Clin Geriatr Med* 1995; 11:159-169.
8. Belchetz PE. Pituitary and adrenal disorders in old age. In: *Textbook of Geriatric Medicine and Gerontology* 4<sup>th</sup> ed. Brocklehurst JC, Tallis RC, Fillit H (eds). Edinburgh, Churchill Livingstone, 1992, pp. 694-700.
9. McMahan DJ, Schwab CW, Kauder D. Comorbidity and the elderly trauma patient. *World J Surg* 1996; 20:1113-1120.
10. Schwab CW, Kauder DR. Geriatric trauma. In: *Early Care of the Injured Patient*. Moore EE, Ducker TB, Edlich FR, et. al. (eds). Toronto, B. D. Decker, 1990, pp. 328-334.
11. Milzman DP, Boulanger BR, Rodriguez A, et. al. Pre-existing disease in trauma patients: a predictor of fate independent of age and ISS. *J Trauma* 1992; 32:236
12. Smith DP, Enderson BL, Maull KI. Trauma in the elderly: determinants of outcome. *South Med J* 1990; 83:171-177.
13. Sartoretti C, Sartoretti-Schefer S, Ruckert R, et. al. Comorbid conditions in old patients with femur fractures. *J Trauma* 1997; 43:570-577.
14. Battistella FD, Din AM, Perez L. Trauma patients 75 years and older: long-term follow-up results justify aggressive management. *J Trauma* 1998; 44:618-624.
15. Sacco WJ, Copes WS, Bain LW, et. al. Effect of preinjury illness on trauma patient survival outcome. *J Trauma* 1993; 35:538-543.
16. MacKenzie EJ, Morris J, Edelstein S. Effect of pre-existing disease on length of hospital stay in trauma patients. *J Trauma* 1989; 29:757-765.
17. Clancy TV, Ramshaw DG, Maxwell JG, et. al. Management outcomes in splenic injury. *Ann Surg* 1997; 226:17-24.
18. DelGuercio LR, Cohn JD. Monitoring operative risk in the elderly. *JAMA* 1980; 243:1350-13
19. Scalea TM, Simon HM, Duncan AO, et al. Geriatric blunt multiple trauma: improved survival with invasive monitoring. *J Trauma* 1990; 30:129-136.

20. Schultz RJ, Whitfield GF, LaMura JJ, et al. The role of physiologic monitoring in patients with fractures of the hip. *J Trauma* 1985; 25:309-316.
21. Tornetta III P, Mostafavi H, Riina, J, et al. Morbidity and mortality in elderly trauma patients. *J Trauma* 1999; 46:702-706.
22. Oreskovich MR, Howard JD, Copass MK, et al. Geriatric trauma: injury patterns and outcome. *J Trauma* 1984; 24:565-572.
23. Perdue PW, Watts DD, Kaufmann CR, et al. Differences in mortality between elderly and younger adult trauma patients: geriatric status increases risk of delayed death. *J Trauma* 1998; 45:805-810.
24. Knudson MM, Lieberman J, Morris JA, et al. Mortality factors in geriatric blunt trauma patients. *Arch Surg* 1994; 129:448-453.
25. Pellicane JV, Byrne K, DeMaria EJ. Preventable complications and death from multiple organ failure among geriatric trauma victims. *J Trauma* 1992; 33:440-444.
26. Horst HM, Obeid FN, Sorenson VJ, Bivins BA. Factors influencing survival of elderly trauma patients. *Crit Care Med* 1986; 14:681-684.
27. Van Aalst JA, Morris JA, Yates HK, et. al. Severely injured geriatric patients return to independent living: a study of factors influencing function and independence. *J Trauma* 1991; 31:1096-1102.
28. Davis JW, Kaups KL. Base deficit in the elderly: a marker of severe injury and death. *J Trauma* 1998; 45:873-877.
29. Demarest GB, Osler TM, Clevenger FW. Injuries in the elderly: evaluation and initial response. *Geriatrics* 1990; 45:36-42.

**Evidentiary Table: Resuscitation Goals in Geriatric Trauma**

<b>First Author</b>	<b>Year</b>	<b>Reference</b>	<b>Data Class</b>	<b># Pts</b>	<b>Age</b>	<b>Pt Population</b>	<b>Mort</b>	<b>Synopsis and Conclusions</b>
Schulz RJ	1985	The role of physiologic monitoring in patients with fractures of the hip. <i>J Trauma</i> ; 25:309-16.	I	35	Mean = 67 in controls	All patients with hip fractures	29% in controls 2.9% in study group	All patients randomly assigned (not blinded) to monitored group or unmonitored group. Hemodynamic monitoring was accomplished using a pulmonary artery catheter. Preoperative risk factors, length of procedure, and postoperative morbidity were similar. The condition of each patient was optimized before surgery using diuretics or inotropes (no formal protocol given). Mortality in monitored group was one-tenth of unmonitored group.
Scaela TM	1990	Geriatric blunt multiple trauma: improved survival with early invasive monitoring. <i>J Trauma</i> ; 30:129-36.	II	1986 – 15 1987 – 30	> 65	Blunt multiple trauma	1986 – 93% 1987 – 47%	1986 Group I 3.5 L/min CO 100% (all cardiogenic shock) Group II 3.5 – 5 86% 1987 Group A 3.5 54% (3 – cardiogenic shock, 4 – organ failure) Group B 3.5 – 5 50% (3 – head injury, 1 – sudden death) Group C > 5 33%
Oreskovich MR	1984	Geriatric trauma: injury patterns and outcome. <i>J Trauma</i> ; 24:565-72.	III	100	> 70	“Severe” blunt trauma; burns included	15%	Attempts were made to optimize to a cardiac index ? 4 L/min/M <sup>2</sup> and/or an oxygen consumption index of 170 cc/min/M <sup>2</sup> . Fluid and inotropes used as needed. Although stable by usual clinical criteria, there may be a dangerous low-flow state. The ability to correct this low-flow state correlates with survival.
Horst HM	1986	Factors influencing survival of elderly trauma patients. <i>Crit Care Med</i> ; 14:681-4.	III	39	? 60	Admitted to SICU. Monitored with PA and arterial catheters.	31%	All non-survivors were in shock (systolic blood pressure < 80 mm Hg) for at least 15 minutes between injury and admission. Only 6% of survivors were found to be in shock. During this study the protocol for prehospital care in the hypotensive patient called for 2,200 ml. lactated Ringer’s solution prior to arrival at hospital.
Pellucane JV	1992	Preventable complications and death from multiple organ failure among geriatric trauma victims. <i>J Trauma</i> ; 33:440-4.	III	374	> 65	Consecutive trauma patients; Burns excluded	8%	Fifteen (38%) of 39 patients presented in shock. Survival related to sepsis and the number of failed organ systems, but NOT presence of shock at admission. Incidence of shock not statistically different between survivors and nonsurvivors. Although survivors had higher mean arterial pressure, cardiac index, left ventricular stroke work, oxygen delivery, and hemoglobin, this was not statistically significant.
Knudson MM	1994	Mortality factors in geriatric blunt trauma	III	852	? 65	Blunt trauma	18.4%	Trauma score was significantly (p < 0.001) higher in patients who survived. Mortality was significantly (p < 0.05) increased in patients with TS < 12 (65%) and TS = 12-14 (25%) when compared with patients with TS = 15-16 (5%). Geriatric patients with a TS < 15 are at high risk and should be admitted to the ICU and treated aggressively.
						Admitting physiologic status predictive of mortality. Systolic blood pressure < 90 mm Hg associated with 82% mortality		

**Evidentiary Table: Resuscitation Goals in Geriatric Trauma**

<b>First Author</b>	<b>Year</b>	<b>Reference</b>	<b>Data Class</b>	<b># Pts</b>	<b>Age</b>	<b>Pt Population</b>	<b>Mort</b>	<b>Synopsis and Conclusions</b>
Davis JW	1998	Base deficit in the elderly: a marker of severe injury and death. <i>J Trauma</i> ; 45:873-7.	III	274	≥ 55	“Major trauma patients” Study group compared to cohort of younger patients	Varied with base deficit	rate. Multiple logistic regression used to construct formula to help predict which patients may benefit from aggressive care. Correlated base deficit with mortality. Arterial blood gases obtained within one hour after admission. Higher mortality in elderly with increasing base deficit, despite similar ISS. In patients ≥ 55 years, a BD 2 to ?2 was associated with an 18% mortality; a BD ?3 to ?5 resulted in 23 % mortality; a BD ?6 to ?9 resulted in 60% mortality; a BD < ?10 resulted in 80% mortality. In all categories, mortality was increased for elderly compared to younger cohort. Positive predictive value not different between elderly and young. Negative predictive value of normal BD in young (60%) was greater than elderly (40%). BD < ?6 is particularly ominous in elderly.
Perdue PW	1998	Differences in mortality between elderly and younger adult trauma patients: geriatric status increases risk of delayed death. <i>J Trauma</i> ; 45:805-10.	III	448	≥ 65	One-system injuries and admits to non-trauma service excluded	14%	Elderly mortality significantly (p < 0.001) greater than that of younger patients. ISS and RTS independently predictive of mortality. Authors practice is to admit elderly patients to ICU if they have significant injury (AIS > 3), shock, or significant chronic cardiovascular or renal disease. Pulmonary artery catheters not placed unless volume or cardiac status uncertain.
Tometta P	1999	Morbidity and mortality in elderly trauma patients. <i>J Trauma</i> ; 46:702-6.	III	326 (Multi-center)	≥ 60	Significant blunt trauma only. Slip-and-fall injuries were excluded	18.1%	Patients who died displayed greater transfusion requirement (10.9 vs. 2.9 units) and more fluid infused (12.4 vs. 4.9 liters). Transfusion requirement and fluid requirement found to be predictive of mortality. Risks of invasive monitoring easily justified in patients with ISS > 18. In patients with ISS < 18, indications need to be evaluated further.