

## Breaking Old Dogma: Continuation of Enteral Feeds up to the Time of Tracheostomy and PEG Placement

### 1. Specific Aims

- a. To determine if continuation of enteral feeds up to the time of tracheostomy improves nutritional status as evident by total caloric intake, and has no effect on risk of aspiration as evident by reported feed material in mouth, worsening consolidation or opacification on chest x-ray within 48 hours, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, and pneumonia within 7 days
- b. To determine if continuation of enteral feeds up to the time of percutaneous endoscopic gastrostomy tube (PEG) placement improves nutritional status as evident by total caloric intake, and has no effect risk of aspiration as evident by reported feed material in mouth, worsening consolidation or opacification on chest x-ray within 48 hours, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, and pneumonia within 7 days

### ***Hypothesis***

We hypothesize that continuing enteral tube feeds in critically ill intubated patients up to the time of both tracheostomy and PEG does not increase risk of aspiration and may improve overall nutritional status. We also hypothesize that there is no difference in aspiration events between nasogastric and nasoduodenal enteral feeding routes or operations performed in the operating room versus in the ICU.

### ***Statement on Impact***

There are currently no standardized guidelines regarding whether intubated patients can continue enteral feeds up to the time of tracheostomy and PEG placement. Additionally, it is unclear whether the route (i.e. nasogastric versus nasoduodenal tube feeds) or the location of the procedure (OR vs ICU) has any implications on feasibility and safety. The purpose of this study is to implement a protocol designed to decrease the interruption of enteral feeds in critically ill patients in hopes of increasing quantitative nutritional intake and prospectively evaluate whether with feeds there is an increased risk of aspiration. We will also use our data to explore the safety and efficacy of nasogastric versus nasoduodenal feeds, and explore differences in subsets of patients to evaluate if there is a specific diagnosis or injury pattern which benefits from continuing feeds compared to others.

Currently at our institution holding feeds preoperatively (6 hours) for tracheostomy is physician dependent while holding feeds for PEG tube placement is standard. In our recent quality improvement retrospective project, we demonstrated that patients who had feeds continued to the time of tracheostomy had no difference in pulmonary complications. We also noted that patients with feeds continued were more likely to have sacral decubitus wounds. As preoperative tube feed management is based on surgeon preference, we suspect this result is due to selection bias, in that the surgeon is more willing to continue feeds in sicker, deconditioned patients with poor nutritional status. We believe that continuing feeds up to the time of tracheostomy and PEG placement will decrease feed interruptions thereby increasing quantitative nutritional intake and show no difference in aspiration rates as defined by feed material in the mouth, worsening consolidation or opacification on chest x-ray within 48 hours post operatively, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, pneumonia within 7 days. Overall, we feel these benefits will ultimately lead to decreases in morbidity associated with malnutrition and will also decrease delays to the operating room.

### 2. Research Strategy

#### **A. Significance:**

Malnutrition is a significant problem in the critically ill, however with busy operating room schedules, we often find that feeds are interrupted repetitively prior to elective surgery. These interruptions are practiced based on the American Society of Anesthesiologist guidelines for fasting intervals for elective surgery. These guidelines, however, were not created for critically ill, intubated patients (14). Controversy around continuing feeds in intubated patients up to the time of surgery exists due to the theoretical concern for loss of airway and need for reintubation on a “full” stomach, and the consequent aspiration risk. Critically ill patients receiving enteral feeds are at risk for aspiration events at baseline with aspiration pneumonitis and aspiration pneumonia resulting in significant morbidity (17). The theoretical concern for an increased risk of aspiration with continuation of enteral feeds up to the time of surgery persists without supporting evidence (1,2,9,10). Not only is ‘to feed or not to feed’ controversial, one study

by Schneider et al demonstrated that anesthesiologist, anesthesia critical care, medical critical care, and surgical critical care physicians often disagree about the length of time necessary to cease the enteral feeding of an intubated patients before scheduled procedures such as tracheostomy (8). It is equally unclear whether nasogastric versus nasoduodenal feeds differ in their ability to administer adequate nutrition and risk of morbidity (11,12,13). Again pointing to the need for a universal evidence based guideline on perioperative enteral feeding to benefit critical care patients.

Additionally, extensive evidence demonstrates that malnutrition is harmful and associated with increased morbidity including increased ICU length of stay and overall increased hospital mortality (15, 16). Meanwhile, good nutrition in the hospital setting is associated with decreased infectious complications, reduced inflammatory cytokines, and improved wound healing (2).

This important topic has been looked at previously by small nonrandomized trials. Angotti et al compared 11 patients who underwent tracheostomy without tube feed interruption to 45 patients with feeds held 4 hours prior to tracheostomy and discovered no difference in pulmonary complications however noted the nil per os group sustained a substantial caloric deficit (10). Pousman et al performed a prospective observational cohort study to evaluate the feasibility of implementing a reduced enteral fasting protocol for mechanically ventilated patients and although no statistical difference was shown, there was a trend demonstrating increased delivery of enteral nutrition and faster achievement of nutrition goals in the intervention group (9). Ho et al developed a protocol for intubated patients undergoing PEG tube placement that involves patients being fed up to the time of the procedure at which point the nasogastric tube is turned to suction to remove residual feeds. The nasogastric tube is continued to suction until the esophagus is intubated by the endoscope at which point the nasogastric tube is removed and the PEG is placed. This group found no periprocedural episodes of vomiting or aspiration, and also noted the stomach had minimal gastric contents upon visualization (2).

To this end, the goal of the present study is to implement a protocol designed to decrease the interruption of enteral feeds in critical care patients and prospectively evaluate whether this leads to a quantitative increase in nutritional intake without increasing the risk of aspiration as defined by reported feed material in mouth, worsening consolidation or opacification on chest x-ray within 48 hours postoperatively, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, and diagnosis of pneumonia within 7 days. Additionally, we hope that increasing quantitative nutrition may decrease morbidity especially in the small subset of patients where feed interruptions occur repetitively.

If the aims of the project are achieved, the data obtained from this study will be used to create an evidenced based platform from which physicians can practice. This will end the anecdotal controversy regarding perioperative tube feed management, increasing quantitative nutrition, and ultimately improving the care of critically ill patients.

## **B. Innovation:**

This project aims to challenge and shift the current clinical practice paradigm of interrupting enteral feeding prior to commonly performed procedures for critical care patients, the tracheostomy and PEG. We hope that through evidence-based medicine we can improve patient care and decrease the risks of malnutrition in the intensive care setting.

## **C. Approach:**

### ***Study design***

To test our hypothesis, we will prospectively identify patients with critical illness requiring tracheostomy and/or PEG after admission to the ICU. Overall patient prognosis and indication for tracheostomy and/or PEG will be determined by the admitting surgeon and the critical care physician. Patients with plans for operative intervention (trach and/or PEG), will be screened and informed consent (via healthcare proxy or next of kin) will be obtained for enrollment.

These patients will then be randomized to our continued enteral feeds arm up to the time of surgery versus nil per os six hours prior. If randomized to the continued enteral feed arm, the patients will continue to receive enteral feeds via their nasogastric or nasoduodenal tube which was placed at the discretion of the critical care physician. Our goal

is to continue tube feeds up to the time of surgery at which time the nasoduodenal tube would be retracted into the stomach and the gastric contents suctioned, or the nasogastric tube is simply put to suction.

We will prospectively follow the patients and evaluate their clinical outcomes including quantitative nutritional benefits and specific indicators for aspiration postoperatively including reported feed material in mouth in the anesthesia record, worsening consolidation or opacification on chest x-ray within 48 hours, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, and diagnosis of pneumonia within 7 days.

We will also collect data, including: diagnosis, medical comorbidities, type of enteral feed, total hours nutrition is held preoperatively, type of nutritional access (NG vs ND), development of ventilator associated pneumonia (VAP) within 7 days, development of surgical site infection, presence of decubitus wounds, and any adverse post-tracheostomy events and/or post percutaneous endoscopic gastrostomy tube events. Data will then be compared in the groups.

In our statistical analysis we will also evaluate for differences between patients receiving feeds via nasogastric tube and nasoduodenal tube up to the time of the procedure. Additionally, will perform subgroup analyses of patients with different diagnosis and injury patterns, and a subgroup analysis on procedures performed in the operating room versus ICU.

### ***Study Population***

Our study population will be adult patients admitted to Surgical and Trauma Intensive Care Units who require tracheostomy and/or PEG tube placement. Retrospective studies of our trauma registry suggest at our single level one trauma center we have 50-80 qualifying patients annually. Once admitted to the ICU, evaluated, and determined to meet inclusion criteria, informed consent for enrollment into the study group will be sought. Given the severity of illness in this patient population, we expect to require the consent of the patient's health care proxy or next of kin.

### ***Inclusion and Exclusion Criteria***

Our study population will be patients admitted to the surgical or trauma ICU who are 18 years or older and require tracheostomy and/ or percutaneous endoscopic gastrostomy tube placement. Patients in whom we are unable to identify a health care proxy or next of kin to obtain informed consent and individuals who are less than 18 years of age will be excluded from this study. Those deemed to be clinically brain dead within 48 hours of enrollment or transitioned to comfort measures within 48 hours of enrollment will also be excluded. Pregnant patients and prisoners will also be excluded from the study.

### ***Data Collected***

Initial data collected upon enrollment will include relevant demographic data including comorbidities and admitting diagnosis. Recent diagnosis of pneumonia (VAP vs CAP) within 7 days of the procedure will also be recorded. Data will then be collected and will include route of enteral access, enteral feed type, enteral feed administration rate, hours enteral feeds help perioperatively, pre-operative and post-operative FIO<sub>2</sub>, preoperative and postoperative ventilator settings, ABG, and new infiltrate on chest x-ray within 48 hours post operatively. We will also review the anesthesia note to evaluate if any aspiration events occurred intraoperatively as evident by suctioning out feeds from the mouth. We will also conduct surveillance for any tracheostomy or PEG related adverse events such as return to OR and surgical site infection.

### ***Study Outcomes***

Quantitative nutritional intake, aspiration events by reported feed material in mouth, worsening consolidation or opacification on chest x-ray within 48 hours, increased requirement of O<sub>2</sub>/ PEEP, worsening ABG, and diagnosis of pneumonia within 7 days.

### ***Study Timeline***

We anticipate the duration of enrollment of this study to last approximately 12 months. Erie County Medical Center is a large, urban Level 1 American College of Surgeons-verified trauma center where we see approximately 50-80 patients per year who meet inclusion criteria. We anticipate in order to have enough data to create a solid statistical analysis we would need an additional 3 centers. Individual subject participation will comprise the length of their hospital stay.

We estimate that completion of primary analysis of the data from this study will be approximately two months following study closing. Data will be collected throughout the study and analyzed to assure subject safety and identification of adverse events. This will be coordinated between the designated co-investigator at each site with the primary investigator. This time frame will allow evaluation of both primary and secondary study endpoints.

#### ***Data Storage and Confidentiality***

All individually identifiable health information will be de-identified prior to entry of subject data into the study database. We plan to use the AAST data collection tool which is password protected. Printed hard copies of data, if used, will be kept in a locked filing cabinet within the principal or co-investigators locked offices to be determined at each site. At Erie County Medical Center this is located at (David K. Miller, C309), within the Department of Surgery at ECMC. If any breach in confidentiality occurs, the principal or co-investigator will contact the Health Sciences Institutional Review Board immediately at the individual center.

#### ***Statistical Analysis***

Standard descriptive statistics such as means, medians, standard deviations for continuous responses (length of stay, ventilator days) and frequencies for categorical responses (complications status, wound/infection status, feed type) will be estimated overall and within each group. For continuous variables, normality will be assessed using goodness of fit tests and quantile-quantile plots. For normally distributed responses, t-tests will be used to evaluate group differences, with Levene's equality of variance test determining the choice of Satterthwaite's approximation or the standard pooled variance estimator. To evaluate group differences in responses exhibiting evidence of a non-normal underlying distribution the Mann-Whitney U test will be used. Association of categorical responses within treatment groups will be evaluated using the chi-squared goodness of fit test.

To assess differences in rates of aspiration events, a log-rank test for differences in survival curves model will be used. Assuming proportional hazards and non-informative censoring, a cox-model will be used to estimate follow-up hazard ratios for development of aspiration pneumonia between the two groups of interest. This assumption has proven reasonable in past research concerning pneumonia development. Should these assumptions not hold, an accelerated failure time model will be used instead, with odds ratios estimated rather than hazard ratios.

To detect a hazard ratio of 0.6667 for aspiration in patients receiving tracheostomy and or PEG placement with continued enteral feeding up to the time of surgery with a type 1 error rate of 0.05 and a power level of 0.80, we require 99 case/control pairs. Pending study enrollment and creation of a multicenter trial, the study should be sufficiently powered.

Follow-up analysis will evaluate risks of tracheostomy or PEG related adverse events, such as return to OR and surgical site infection. Patients who do and do not experience these adverse events will be matched using propensity score techniques using baseline data. The matched sample and standard sample will be examined using survival analysis methods to identify potential risk factors.

#### ***Provisions to Monitor Data and Ensure Safety of Subjects***

To ensure safety of the research subjects, data collected during the study period will be periodically evaluated. Data regarding the primary endpoint as well as morbidity-specific endpoints will be evaluated throughout the study. Data collection for individual subjects will be an ongoing process throughout the study and will be completed on a weekly basis. The first formal data analysis will occur at two months into the study and then two months thereafter until the conclusion of the study.

The data will primarily be analyzed by the principal investigator. We will evaluate the data as described in subsection of this proposal "*Statistical Analysis*". After analyzing the data, it will be presented to a committee composed of the principal investigator, attending surgeons, and intensivists at each site. At these meetings, in addition to evaluating the primary and secondary endpoints we will also review the incidence of any post-tracheostomy or post-PEG occurrences.

#### ***Risks to Subjects***

The physical risks to subjects in this study are minimal as all interventions are within the standard of care. However, we will be specifically tracking and identifying any adverse tracheostomy or PEG related events, should they occur. While this subject population (critically ill) is at risk for psychological and social/economic risks

following severe illness/injury (PTSD, anxiety, depression, altered behavior, damage to employability, etc.), we do not foresee any direct risks due to participation.

### ***Consent Process***

Informed consent for enrollment into this study will be obtained by the primary co-investigator at each site after determining patients meeting inclusion criteria. The principal investigator at each site will be a member of the surgical or intensive care team and will be involved in the consent process. The consent process will take place after determining the patient's appropriateness for the study. This process will be completed in the intensive care unit.

The consent process will be extensively discussed with the patient's health care proxy or next of kin to ensure an understanding of the study and those consenting will be asked to expressly recite their understanding of the process. The documentation of informed consent in writing will be completed with a member of the nursing staff in attendance who will attest to the consent process.

We hypothesize that continuing enteral feeds in critically ill intubated patients up to the time of tracheostomy and PEG tube placement increases overall quantitative nutritional intake, has no increase in aspiration risk, and decreases morbidity. We also hypothesize no difference in morbidity between nasogastric and nasoduodenal enteral feeding routes and no difference in procedures performed in the operating room compared to in the intensive care setting. We also hypothesize a benefit in feeding with isolated head or extremity pathology compared to those with abdominal pathology. Our goal is to be able to implement a new standard of care designed to decrease the interruption of enteral feeds in critical care patients thereby increasing nutritional intake and decreasing morbidity.

### ***Data Dictionary***

Pneumonia- Chest XRAY with new or progressive infiltrate, consolidation, cavitation AND either fever  $>38^{\circ}\text{C}$  or leukopenia ( $<4000$  wbc) or leukocytosis ( $>12,000$  wbc) , AND new onset purulent sputum, increased secretions, increased suctioning requirement, new onset tachypnea, dyspnea or cough, or increased oxygen requirements, ventilator demands or worsening gas exchange

VAP (ventilator associated pneumonia)- The following clinical signs and symptoms after  $\geq 2$  days of mechanical ventilation: Chest XRAY with new or progressive infiltrate, consolidation, cavitation AND either fever  $>38^{\circ}\text{C}$  or leukopenia ( $<4000$  wbc) or leukocytosis ( $>12,000$  wbc) , AND new onset purulent sputum, increased secretions, increased suctioning requirement, new onset tachypnea, dyspnea or cough, or increased oxygen requirements, ventilator demands or worsening gas exchange

SSI (Surgical Site Infection)- Infection which occurs after surgery in the part of the body where the surgery took place

Immediate adverse post-operative event- death within 24 hours of surgery, myocardial infarction within 24 hours, stroke within 24 hours, gastric contents within mouth/ emesis within 24 hours, bleeding from surgical site within 24 hours, reoperation within 24 hours

### ***Bibliography:***

1. G E O'Keefe et al. Safety of minimizing preoperative starvation in critically ill and intubated trauma patients. *Journal of Trauma and Acute Care Surgery* 2015.
2. Ho C, Culhane J. Reduced Fasting Protocol for Endoscopic Percutaneous Gastrostomy in Intubated Patients. *Int J Clin Med*. 2013;4(8):369-373.
3. K K Singh et al An integrated systematic and meta-analysis of published randomized controlled trials evaluating nasogastric against postpyloris (nasoduodenal and nasojejunal) feeding in critically ill patients admitted to the intensive care unit. *European Journal of Clinical Nutrition* 2014
4. Kortbeek JB, Haigh PI, Doig C. Duodenal versus gastric feeding in ventilated blunt trauma patients: a randomized controlled trial. *J Trauma*. 1999;46:992-998.
5. Nutritional Support: Site of enteral support Gastric versus Jejunal. EAST Guidelines. *J Trauma* September 2004
6. R Fabia et al. Safety and efficacy of uninterrupted perioperative enteral feeding in pediatric burn patients. *Burns* 2018

7. Richardson et al. Transpyloric Passage of Feeding Tubes in Patients with Head Injuries Does Not Decrease Complications. *Journal of Trauma: Injury, Infection, and Critical Care* 1995
8. Schneider et al. Institutional practices of withholding enteral feeding from intubated patients. *Journal of Critical Care Medicine* 7. 2009
9. R. M. Pousman, C. Pepper, P. Pandharipande, G. D. Ayers, B. Mills, J. Diaz, B. Collier, R. Miller and G. Jensen, "Feasibility of Implementing a Reduced Fasting Protocol for Critically Ill Trauma Patients Undergoing Operative and Nonoperative Procedures," *Journal of Parenteral and Enteral Nutrition*, Vol. 33, No. 2, 2009, pp. 176-180
10. L. Angotti et al. New Protocol Avoiding Tube Feed Interruptions in Critically Ill Patients Requiring Tracheostomy. *AM Surg.* 2018
11. Sajid et al. An Integrated Systematic Review and Meta-Analysis of Published Randomized Controlled Trials Evaluating Nasogastric against Postpylorus Feeding in Critically Ill Patients Admitting in Intensive Care Units. *Eur J Clin Nutrition.* 2014 Apr;68(4):424-32.
12. Duodenal Versus Gastric Feeding in Medical Intensive Care Unit Patients: A Prospective Randomized, Clinical Study. *Critical Care Med* 2009 Jun;37(6):1866-72.
13. P Marik et al. Gastric Versus Post-Pyloric Feeding: A Systematic Review. *Critical Care.* 2003 Jun;7(3):R46-51.
14. Practice Guidelines for Preoperative Fasting and the Use of Pharmacologic Agents to Reduce the Risk of Pulmonary Aspiration: Application to Healthy Patients Undergoing Elective Procedures. An Updated Report by The American Society of Anesthesiologists Task Force on Preoperative Fasting and the Use of Pharmacologic Agents to Reduce the Risk of Pulmonary Aspiration. *Anesthesiology* Mar 2017.
15. Han et al. Association Between Malnutrition and Clinical Outcomes in the Intensive Care Unit: A Systematic Review. *Journal of Parenteral and Enteral Nutrition.* Feb 2016
16. Havens et al. Malnutrition at Intensive Care Unit Admission Predicts Mortality in Emergency General Surgery Patients. *JPEN.* 2018 Jan;42(1):156-163.
17. DiBardino et al. Aspiration pneumonia: A review of modern trends. *J Critical Care.* Feb 2015
18. Pneumonia (Ventilator-associated [VAP] and nonventilator-associated Pneumonia [PNEU]) Event. [Pneumonia \(cdc.gov\)](https://www.cdc.gov/pneumonia/)